TWIN-WIDTH AND PERMUTATIONS

ÉDOUARD BONNET $@~^a,$ JAROSLAV NEŠETŘIL $@~^b,$ PATRICE OSSONA DE MENDEZ $@~^{c,b},$ SEBASTIAN SIEBERTZ $@~^d,$ AND STÉPHAN THOMASSÉ $@~^a$

^a Université de Lyon, CNRS, ENS de Lyon, France e-mail address: edouard.bonnet@ens-lyon.fr, stephan.thomasse@ens-lyon.fr

^b Computer Science Institute of Charles University (IUUK), Praha, Czech Republic *e-mail address*: nesetril@iuuk.mff.cuni.cz

 c Centre d'Analyse et de Mathématiques Sociales, EHESS, CNRS UMR 8557, Paris, Francee-mail address: pom@ehess.fr

^d University of Bremen, Bremen, Germany *e-mail address*: siebertz@uni-bremen.de

> ABSTRACT. Inspired by a width invariant on permutations defined by Guillemot and Marx, Bonnet, Kim, Thomassé, and Watrigant introduced the twin-width of graphs, which is a parameter describing its structural complexity. This invariant has been further extended to binary structures, in several (basically equivalent) ways. We prove that a class of binary relational structures (that is: edge-colored partially directed graphs) has bounded twin-width if and only if it is a first-order transduction of a proper permutation class. As a by-product, we show that every class with bounded twin-width contains at most $2^{O(n)}$ pairwise non-isomorphic *n*-vertex graphs.

1. INTRODUCTION

In this paper we consider the graph parameter *twin-width*, defined by Bonnet, Kim, Thomassé and Watrigant [BKTW22] as a generalization of an invariant for classes of permutations defined by Guillemot and Marx [GM14]. Twin-width was recently studied intensively in the context of many structural and algorithmic questions, such as FPT model checking [BKTW22], graph enumeration [BGK⁺21a], graph coloring [BGK⁺21b], and structural properties of matrices and ordered graphs [BGO⁺24].

This paper is part of a project that has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (grant agreement No 810115 – DYNASNET) and from the German Research Foundation (DFG) with grant agreement No 444419611.





DOI:10.46298/LMCS-20(3:4)2024

Key words and phrases: Twin-width, first-order transductions, structural graph theory.

Many well-studied classes of graphs have bounded twin-width: planar graphs, and more generally, any class of graphs excluding a fixed minor, cographs, and more generally, any class of bounded clique-width, etc.

The twin-width of graphs was originally defined using a sequence of 'near-twin' vertex contractions or identifications. Roughly speaking, twin-width measures the accumulated error (recorded via the so-called 'red edges') made by the identifications. To help the reader start forming intuitions, we give a concise definition of the twin-width of a graph; a formal generalization for binary structures is presented in Section 2.4.

A trigraph is a graph with some edges colored red (while the rest of them are black). A contraction (or identification) consists of merging two (non-necessarily adjacent) vertices, say, u, v into a vertex w that is adjacent to a vertex z via a black edge if uz and vz were black edges, or otherwise, via a red edge if at least one of u and v were adjacent to z. The rest of the trigraph does not change. A contraction sequence of an n-vertex graph G is a sequence of trigraphs $G = G_n, \ldots, G_1$ such that G_i is obtained from G_{i+1} by performing one contraction (observe that G_1 is the 1-vertex graph). A *d*-sequence is a contraction sequence where all the trigraphs have red degree at most d. The twin-width of G is then the minimum integer d such that G admits a d-sequence. See Figure 1 for an example of a graph admitting a 2-sequence.




Figure 1: A 2-sequence witnessing that the initial graph has twin-width at most 2.

In this paper, the extension of twin-width for binary relational structures perfectly matches the one in [BKTW22] on undirected graphs, but will slightly differ for general binary structures. Though, as we will observe, both definitions give parameters that differ only by, at most, a linear factor.

We show that twin-width can be concisely expressed by special structures, which we call *twin-models*. Twin-models are rooted trees augmented by a set of transversal edges that satisfies two simple properties: minimality and consistency. These properties imply that every twin-model admits a *ranking*, from which we can compute a *width*. The twin-width of a structure then coincides with the optimal width of a ranked twin-model of the structure. While this connection is technical, twin-models provide a simple way to handle classes of binary structures with bounded twin-width. Note that an informal precursor of ranked twin-models appears in [BGK⁺21b] in the form of the so-called *ordered union trees* and the realization that the edge set of graphs of twin-width at most d can be partitioned into $O_d(n)$ bicliques where both sides of each biclique are a discrete interval along a unique fixed vertex ordering. The main novelty in the (ranked) twin-models lies in the axiomatization of *legal* sets of transversal edges, which is indispensable to their logical treatment.

This paper is a combination of model-theoretic tools (relational structures, interpretations, transductions), structural graph theory and theory of permutations. Here, by a permutation, we mean a relational structure consisting of two linear orders on the same set (see [ABF20]

for a discussion on representations of permutations). Note that this type of representation is particularly adapted to the study of patterns in permutations. The following is the main result of this paper:

Theorem. A class of binary relational structures has bounded twin-width if and only if it is a first-order transduction of a proper permutation class.

The "only if" part of this theorem is stated in more technical terms in Section 7 as Theorem 7.2, and is our main contribution. The other direction, the fact that every binary structure that is a first-order transduction of a proper permutation class has bounded twin-width, was already known [BKTW22]. More specifically, it was shown that proper permutation classes have bounded twin-width [BKTW22, Section 6.1] and that every firstorder transduction of a class of bounded twin-width has itself bounded twin-width [BKTW22, Theorem 8.1].

We recall that a proper permutation class is a set of permutations closed under subpermutations that excludes at least one permutation. Transductions provide a model theoretical tool to encode relational structures (or classes of relational structures) inside other (classes of) relational structures and will be formally defined in Section 2.3.

The fact that any class of graphs with bounded twin-width is just a transduction of a very simple class (a proper permutation class) is surprising at first glance, and it nicely complements another model theoretic characterization of classes of bounded twin-width: a class of graphs has bounded twin-width if and only if it is the reduct of a dependent class of ordered graphs [BGO⁺24]. It can also be thought of as scaling up the fact that classes of bounded rank-width coincide with transductions of tree orders, and classes of bounded linear rank-width, with transductions of linear orders [Col07]. On the other hand, twin-models are interesting objects per se and in a way present one of the most permissive forms of width parameters related to trees. Note that for other classes of sparse structures we do not have such concrete models.

The main result implies that every relational structure on n elements from a class with bounded twin-width can be encoded in a permutation on at most kn elements for some number k. It is then a consequence of [MT04] that every class of relational structures with bounded twin-width contains at most c^n non-isomorphic structures with n vertices, hence is small (i.e., contains at most $c^n n!$ labeled structures with n elements). This extends the main result of [BGK⁺21a] while not using the "versatile twin-width" machinery (but only the preservation of bounded twin-width by transductions proved in [BKTW22]). This also extends a similar property for proper minor-closed classes of graphs, which can be derived from the boundedness of book thickness, as noticed by McDiarmid (see the concluding remarks of [BNW10]).

The proof of our main result is surprisingly complex and proceeds in several steps, which perhaps add new aspects to the rich spectrum of structures related to twin-width. The basic steps can be outlined as follows (the relevant terminology will be formally introduced in the appropriate sections).

We start with a class \mathscr{C}_0 of binary relational structures with bounded twin-width. We derive a class \mathscr{T} of twin-models (tree-like representations of the structures using rooted binary trees and transversal binary relations). Replacing the rooted binary trees of the twin-models by binary tree orders, we get a class \mathscr{F} of so-called full twin-models, which we prove has bounded twin-width. This class can be used to retrieve \mathscr{C}_0 as a transduction, that is by means of a logical encoding. Using a transduction pairing (generalizing the notion of a bijective encoding) between binary tree orders \mathscr{O} and rooted binary trees ordered by

Vol. 20:3

a preorder $\mathscr{Y}^{<}$ we derive a transduction pairing of the class of full twin-models \mathscr{F} with a class $\mathscr{T}^{<}$ of ordered twin-models. From the property that the class \mathscr{G} of the Gaifman graphs of the twin-models in \mathscr{T} is degenerate (and has bounded twin-width), we prove a transduction pairing of \mathscr{T} and \mathscr{G} , from which we derive a transduction pairing of $\mathscr{T}^{<}$ and the class $\mathscr{G}^{<}$ of ordered Gaifman graphs of the ordered twin-models. As a composition of a transduction pairing of $\mathscr{G}^{<}$ with a class $\mathscr{E}^{<}$ of ordered binary structures, in which each binary relation induces a pseudoforest and a transduction pairing of $\mathscr{E}^{<}$ with a class \mathscr{P} of permutations we define a transduction pairing of $\mathscr{G}^{<}$ and \mathscr{P} . As $\mathscr{G}^{<}$ has bounded twin-width (as it is a transduction of a class with bounded twin-width) we infer that \mathscr{P} avoids at least one pattern. Following the backward transductions, we eventually deduce that \mathscr{C}_{0} is a transduction of the hereditary closure $\overline{\mathscr{P}}$ of \mathscr{P} , which is a proper permutation class.

This proof may be schematically outlined by Figure 2. Here, all the notations are consistent with the notation used later in our proof.




Figure 2: Relations between the classes of structures involved in the proof of the main result. The interpretation S is defined in Definition 5.1, the transduction pairing (L, O) in Lemma 6.1, the transduction pairing $(\widehat{L}, \widehat{O})$ as a remark just after Definition 6.2, the transduction pairing (G, U) in Lemma 6.4, and the transduction pairing $(\widehat{G}, \widehat{U})$ as a remark just after Definition 6.5.

The full transformation of a graph G into a permutation σ and the inverse transformation (obtained as a transduction) are displayed on Figure 3 on an example.




Figure 3: From a graph G to a permutation σ , and back.

2. Preliminaries

2.1. Relational structures. We assume basic knowledge of first-order logic and refer to [H⁺97] for extensive background. A relational signature Σ is a finite set of relation symbols R_i with associated arity r_i . A relational structure **A** with signature Σ , or simply a Σ -structure consists of a domain A together with relations $R_i(\mathbf{A}) \subseteq A^{r_i}$ for each relation symbol $R_i \in \Sigma$ with arity r_i . The relation $R_i(\mathbf{A})$ is called the *interpretation* of R_i in **A**. We will often speak of a relation instead of a relation symbol when there is no ambiguity. We may write **A** as $(A, R_1(\mathbf{A}), \ldots, R_s(\mathbf{A}))$. In this paper we will consider relational structures with finite We will further assume that Σ -structures are *irreflexive*, that is, $(v, v) \notin R_i(\mathbf{A})$ for every element $v \in A$ and relation symbol $R_i \in \Sigma$. A unary relation is called a *mark*. Let R be a binary relation symbol and let $u, v \in A$. That the pair (u, v) lies in the

Vol. 20:3

interpretation of R in \mathbf{A} will be indifferently denoted by $(u, v) \in R(\mathbf{A})$ or $\mathbf{A} \models R(u, v)$. More generally, for a formula $\varphi(x_1, \ldots, x_k)$, a Σ -structure \mathbf{A} , an integer $\ell < k$ and $a_1, \ldots, a_\ell \in A$ we define $\varphi(\mathbf{A}, a_1, \ldots, a_\ell) := \{(x_1, \ldots, x_{k-\ell}) \in A^{k-\ell} : \mathbf{A} \models \varphi(x_1, \ldots, x_{k-\ell}, a_1, \ldots, a_\ell)\}$. In this paper, by formula, we mean a first-order formula in the language of Σ -structures, where Σ is usually understood from the context. Let $\mathbf{A} = (A, R_1(\mathbf{A}), \ldots, R_s(\mathbf{A}))$ be a Σ -structure and let $X \subseteq A$. The substructure of \mathbf{A} induced by X is the Σ -structure $\mathbf{A}[X] = (X, R_1(\mathbf{A}) \cap X^{r_1}, \ldots, R_k(\mathbf{A}) \cap X^{r_s})$.

Graphs are structures with a single binary relation E encoding adjacency; this relation is irreflexive and symmetric. Graphs of particular interest in this paper are rooted trees. For a rooted tree Y, we denote by I(Y) the set of internal nodes of Y, by L(Y), the set of leaves of Y, by $V(Y) = I(Y) \cup L(Y)$ the set of vertices of Y, by r(Y), the root of Y, and by \preceq_Y , the partial order on V(Y) defined by $u \preceq_Y v$ if the unique path in Y linking r(Y)and v contains u (i.e., if u = v or u is an ancestor of v in Y). For a non-root vertex v, we further denote by $\pi_Y(v)$ the parent of v, which is the unique neighbor of v smaller than vwith respect to \preceq_Y . (We further define $\pi_Y(r(Y)) = r(Y)$, so that π_Y is defined on all the vertices of Y, the root being the only fixed point.) A rooted binary tree is a rooted tree such that every internal node has exactly two children.

Let Y be a rooted tree and let A be a subset of vertices of Y closed by pairwise least common ancestor (that is: the least common ancestor in Y of any two vertices in A also belongs to A). The subtree of Y induced by A is the rooted tree Y', whose associated tree order $\leq_{Y'}$ is the restriction to A of the tree order \leq_Y associated to Y. In particular, A is the vertex set of Y'.

Partial orders are structures with a single antisymmetric and transitive binary relation \prec . Particular partial orders will be of interest here. Linear orders (also called total orders) are partial orders such that $\forall x \forall y ((x \prec y) \lor (y \prec x) \lor (y = x))$. Tree orders are partial orders that satisfy the following axioms: $\forall x \forall y \forall z ((x \prec z \land y \prec z) \rightarrow ((x \prec y) \lor (y \prec x) \lor (x = y)))$ and $\exists r \forall x ((x = r) \lor (r \prec x))$. The minimum element of a tree order (r in the previous equation) is its root, and its maximal elements are its leaves. It will be convenient to use \preceq, \succ, \succeq with their obvious meaning. Let (X, \prec) be a tree order. The infimum inf(u, v) of two elements $u, v \in X$ is the unique element $w \in X$ such that $w \preceq u, w \preceq v$, and $\forall z (((z \preceq u) \land (z \preceq v)) \rightarrow (z \preceq w)))$. Note that $\inf(x, y)$ is first-order definable from \prec , hence can be used as a term in our formulas. An element x is covered by an element y if $x \prec y$ and there is no element z with $x \prec z \prec y$. A binary tree order is a tree order such that every non-maximal element is covered by exactly two elements.

Ordered graphs are structures with two binary relations, E and <, where E defines a graph and < defines a linear order. We denote ordered graphs as $G^{<} = (V, E, <)$.

A permutation is represented as a structure $\sigma = (V, <_1, <_2)$, where V is a finite set and where $<_1$ and $<_2$ are two linear orders on this set (see e.g. [Cam02, ABF20]). Two permutations $\sigma = (V, <_1, <_2)$ and $\sigma' = (V', <'_1, <'_2)$ are isomorphic if there is a bijection between V and V' preserving both linear orders. Let $X \subseteq V$. The sub-permutation of σ induced by X is the permutation on X defined by the two linear orders of σ restricted to X. The isomorphism types of the sub-permutations of a permutation σ are the patterns of σ . A class \mathscr{P} of (isomorphism types of) permutations is hereditary (or closed) if it is closed under taking sub-permutations. A permutation class is a hereditary class of permutations. A permutation class is proper if it is not the class of all permutations. Note that the terms "class of permutations" and "permutation class" are not equivalent, the second referring to a hereditary class of permutations, as it is customary (see e.g. [Bón12]). 2.2. Interpretations. Let Σ, Σ' be signatures. A simple interpretation (or, simply, an interpretation, since we will only consider these in this article) I of Σ' -structures in Σ -structures is defined by a Σ -formula $\rho_0(x)$, and a Σ -formula $\rho_{R'}(x_1, \ldots, x_k)$ for each k-ary relation symbol $R' \in \Sigma'$. Let I be an interpretation of Σ' -structures in Σ -structures, where $\Sigma' = \{R'_1, \ldots, R'_s\}$. For each Σ -structure **A** we denote by $I(\mathbf{A}) = (\rho_0(\mathbf{A}), \rho_{R'_1}(\mathbf{A}), \ldots, \rho_{R'_s}(\mathbf{A}))$ the Σ' -structure interpreted by I in **A**. Similarly, for a class \mathscr{C} of Σ -structures, we denote by $I(\mathscr{C})$ the set $\{I(\mathbf{A}) : \mathbf{A} \in \mathscr{C}\}$.

We denote by $\operatorname{\mathsf{Reduct}}_{\Sigma^+\to\Sigma}$ (or simply $\operatorname{\mathsf{Reduct}}$ when Σ and Σ^+ are clear from context) the interpretation that "forgets" the relations in $\Sigma^+ \setminus \Sigma$ while preserving all the other relations and the domain. For a Σ^+ -structure \mathbf{B} , the Σ -structure $\operatorname{\mathsf{Reduct}}(\mathbf{B})$ is called the Σ -reduct (or simply reduct if Σ is clear from the context) of \mathbf{B} . A class \mathscr{C} is a reduct of a class \mathscr{D} if $\mathscr{C} = \operatorname{\mathsf{Reduct}}(\mathscr{D})$. Conversely, a class \mathscr{D} is an expansion of \mathscr{C} if \mathscr{C} is a reduct of \mathscr{D} .

Another important interpretation is Gaifman_{Σ} (or simply Gaifman when Σ is clear from context), which maps a Σ -structure **A** to its *Gaifman graph*, whose vertex set is A and whose edge set is the set of all pairs of distinct vertices included in a tuple of some relation.

Note that an interpretation of Σ_2 -structures in Σ_1 -structure naturally defines an interpretation of Σ_2^+ -structures in Σ_1^+ -structures if $\Sigma_2^+ \setminus \Sigma_2 = \Sigma_1^+ \setminus \Sigma_1$ by leaving the relations in $\Sigma_1^+ \setminus \Sigma_1$ unchanged (that is, by considering $\rho_R(x_1, \ldots, x_k) = R(x_1, \ldots, x_k)$ for these relations).

2.3. Transductions. Let Σ, Σ' be signatures. A simple transduction T from Σ -structures to Σ' -structures is defined by a simple interpretation I_T of Σ' -structures in Σ^+ -structures, where Σ^+ is a signature obtained from Σ by adding finitely many marks. For a Σ -structure \mathbf{A} , we denote by $T(\mathbf{A})$ the set of all $I_T(\mathbf{B})$ where \mathbf{B} is a Σ^+ -structure with reduct \mathbf{A} : $T(\mathbf{A}) =$ $\{I_T(\mathbf{B}) : \text{Reduct}(\mathbf{B}) = \mathbf{A}\}$. Let $k \in \mathbb{N}$. The k-blowing of a Σ -structure \mathbf{A} is the Σ' -structure $\mathbf{B} = \mathbf{A} \bullet k$, where Σ' is the signature obtained from Σ by adding a new binary relation \sim encoding an equivalence relation. The domain of $\mathbf{A} \bullet k$ is $B = A \times [k]$, and, denoting by p_1 and p_2 the projections $A \times [k] \to A$ and $A \times [k] \to [k]$ we have, for all $x, y \in B$, $\mathbf{B} \models x \sim y$ if $p_1(x) = p_1(y)$, and (for $R \in \Sigma$) $\mathbf{B} \models R(x_1, \ldots, x_s)$ if $\mathbf{A} \models R(p_1(x_1), \ldots, p_1(x_s))$ and $p_2(x_1) = \ldots = p_2(x_s)$. A copying transduction is the composition of a k-blowing and a simple transduction; the integer k is the blowing factor of the copying transduction T and is denoted by bf(T). It is easily checked that the composition of two copying transductions is again a copying transduction. In the following by the term transduction we mean a copying transduction. Note that for every transduction T from Σ -structure to Σ' , for every Σ -structure \mathbf{A} and for every Σ' -structure $\mathbf{B} \in T(\mathbf{A})$ we have $|B| \leq bf(T) |A|$.

Let T, T' be transductions from Σ -structures to Σ' -structures, and let \mathscr{C} be a class of Σ -structures. The transduction T' subsumes the transduction T on \mathscr{C} if $\mathsf{T}'(\mathbf{A}) \supseteq \mathsf{T}(\mathbf{A})$ for all $\mathbf{A} \in \mathscr{C}$. If \mathscr{C} is a class of Σ -structures, we define $\mathsf{T}(\mathscr{C}) = \bigcup_{\mathbf{A} \in \mathscr{C}} \mathsf{T}(\mathbf{A})$. We say that a class \mathscr{D} of Σ' -structures is a T -transduction of \mathscr{C} if $\mathscr{D} \subseteq \mathsf{T}(\mathscr{C})$ and, more generally, the class \mathscr{D} is a transduction of the class \mathscr{C} , and we write $\mathscr{C} \longrightarrow \mathscr{D}$, if there exists a transduction T such that \mathscr{D} is a T -transduction of \mathscr{C} . The negation of $\mathscr{C} \longrightarrow \mathscr{D}$ is denoted by $\mathscr{C} \longrightarrow \mathscr{D}$. Note that we require only the inclusion of \mathscr{D} in $\mathsf{T}(\mathscr{C})$, and not the equality. The class \mathscr{D} is a *c*-bounded T -transduction of the class \mathscr{C} if, for every $\mathbf{B} \in \mathscr{D}$, there exists $\mathbf{A} \in \mathscr{C}$ with $\mathbf{B} \in \mathsf{T}(\mathbf{A})$ and $|\mathcal{A}| \leq c |\mathcal{B}|$. Two classes \mathscr{C} and \mathscr{D} are transduction equivalent if each is a

transduction of the other. A transduction pairing of two classes \mathscr{C} and \mathscr{D} is a pair (D, C) of (copying) transductions, such that $\forall \mathbf{A} \in \mathscr{C} \exists \mathbf{B} \in D(\mathbf{A}) \cap \mathscr{D} : \mathbf{A} \in C(\mathbf{B})$ and $\forall \mathbf{B} \in \mathscr{D} \\ \exists \mathbf{A} \in C(\mathbf{B}) \cap \mathscr{C} : \mathbf{B} \in D(\mathbf{A}).$

We denote by $\mathscr{C} = \mathscr{D}$ the existence of a transduction pairing of \mathscr{C} and \mathscr{D} . Note that if (D, C) is a transduction pairing, then \mathscr{D} is a bf(C)-bounded D-transduction of \mathscr{C} and \mathscr{C} is a bf(D)-bounded C-transduction of \mathscr{D} . The following easy lemma will be useful.

Lemma 2.1. Assume \mathscr{D} is a D-transduction of \mathscr{C} , \mathscr{C} is a C-transduction of \mathscr{D} , and for every $\mathbf{A} \in \mathscr{C}$ and every $\mathbf{B} \in \mathsf{D}(\mathbf{A}) \cap \mathscr{D}$ we have $\mathbf{A} \in \mathsf{C}(\mathbf{B})$. Then (D,C) is a transduction pairing of \mathscr{C} and \mathscr{D} .

Proof. Let $\mathbf{B} \in \mathscr{D}$. As \mathscr{D} is a D-transduction of \mathscr{C} there exists $\mathbf{A} \in \mathscr{C}$ with $\mathbf{B} \in \mathsf{D}(\mathbf{A})$. Then $\mathbf{A} \in \mathsf{C}(\mathbf{B}) \cap \mathscr{C}$.

Note that a transduction T from Σ_1 -structures to Σ_2 -structures naturally defines a transduction $\widehat{\mathsf{T}}$ from Σ_1^+ -structures to Σ_2^+ -structures if $\Sigma_2^+ \setminus \Sigma_2 = \Sigma_1^+ \setminus \Sigma_1$ by leaving the relations in $\Sigma_1^+ \setminus \Sigma_1$ unchanged. The transduction $\widehat{\mathsf{T}}$ is called the *natural generalization* of T to Σ_1^+ -structures.

2.4. **Twin-width.** In order to define twin-width, we first need to introduce some preliminary notions, which generalize the notion of trigraphs (i.e., graphs with some red edges) introduced in [BKTW22]. Let Σ be a binary relational signature. The signature Σ^* is obtained by adding, for each binary relation symbol R a new binary relation symbol R^* . The symbol R^* will always be interpreted as a symmetric relation and plays for R the role of red edges in [BKTW22].

Let **A** be a Σ^* -structure, and let u and v be vertices of **A**. The vertices u, v are R-clones for a vertex w and a relation $R \in \Sigma$ if we have $\mathbf{A} \models (R(u, w) \leftrightarrow R(v, w)) \land (R(w, u) \leftrightarrow R(w, v))$ and no pair in R^* contains both w and either u or v. The Σ^* -structure \mathbf{A}' obtained by contracting u and v into a new vertex z is defined as follows:

- $A' = A \setminus \{u, v\} \cup \{z\};$
- $R(\mathbf{A}') \cap ((A' \setminus \{z\}) \times (A' \setminus \{z\})) = R(\mathbf{A}) \cap ((A \setminus \{u, v\}) \times (A \setminus \{u, v\}))$ for all $R \in \Sigma^*$;
- for every vertex $w \in A' \setminus \{z\}$ and every $R \in \Sigma$ such that u and v are R-clones for w, we let $\mathbf{A}' \models R(w, z)$ if $\mathbf{A} \models R(w, u)$, and $\mathbf{A}' \models R(z, w)$ if $\mathbf{A} \models R(u, w)$. (Note that this does not change if we use v instead of u);
- otherwise, for every vertex $w \in A' \setminus \{z\}$ and every $R \in \Sigma$ such that u and v are not R-clones for w we let $\mathbf{A}' \models R^*(w, z) \land R^*(z, w)$.

A *d*-sequence for a Σ -structure \mathbf{A} is a sequence $\mathbf{A}_n, \ldots, \mathbf{A}_1$ of Σ^* -structures such that: \mathbf{A}_n is isomorphic to \mathbf{A} (when considered as a Σ^* -structure with empty R^*); \mathbf{A}_1 is the Σ^* structure with a single element; for every $1 \leq i < n$, \mathbf{A}_i is obtained from \mathbf{A}_{i+1} by performing a single contraction; for every $1 \leq i < n$ and every $v \in A_i$, the sum of the degrees in relations $R^* \in \Sigma^* \setminus \Sigma$ of v in \mathbf{A}_i is less or equal to d (the degree of v in relation R^* is defined as the degree of v in the undirected graph $(A, R^*(\mathbf{A}))$). When d is not specified, we shall speak of a contraction sequence; see Figure 4 for an illustration. The minimum d such that there exists a d-sequence for a Σ -structure \mathbf{A} is the twin-width tww(\mathbf{A}) of \mathbf{A} . This definition for binary relational structures differs from the one given in [BKTW22] (where red edges are not counted with multiplicity), but will be more convenient in our setting. However, the definitions differ by at most a constant factor (linear in $|\Sigma|$), thus the derived notion of class with bounded twin-width coincides.




Figure 4: A contraction sequence, a so-called block representation of the contractions, and a twin-model.

A crucial property of twin-width is the following result.

Theorem 2.2 [BKTW22, Theorem 8.1]. Let \mathcal{C}, \mathcal{D} be classes of binary structures. If \mathcal{C} has bounded twin-width and \mathcal{D} is a transduction of \mathcal{C} , then \mathcal{D} has bounded twin-width.

3. Classes with bounded star chromatic number

One of the key ingredients of the proof will rely on a transduction pairing between a class \mathscr{C} and the class of Gaifman graphs of the structures in \mathscr{C} . Though such a pairing does not exist for general classes of structures (see the discussion below), we prove in this section that this is the case if the Gaifman graphs have bounded star chromatic number.

Recall that a *star coloring* of a graph G is a proper coloring of G such that any two color classes induce a star forest (i.e., a disjoint union of stars); the *star chromatic number* $\chi_{st}(G)$ of G is the minimum number of colors in a star coloring of G. Note that a star coloring of a graph with c colors defines a partition of the edge set into $\binom{c}{2}$ star forests. Although we are interested only in binary relational structures in this paper, the next lemma holds (and is proved) for general relational signatures.

Lemma 3.1. Let Σ be a relational signature, let \mathscr{C} be a class of Σ -structures, and let c be an integer. There exists a simple transduction $\mathsf{Unfold}_{\Sigma,c}$ from graphs to Σ -structures such that if the Gaifman graphs of the structures in \mathscr{C} have star chromatic number at most c, then $(\mathsf{Gaifman}_{\Sigma}, \mathsf{Unfold}_{\Sigma,c})$ is a transduction pairing of $(\mathscr{C}, \mathsf{Gaifman}_{\Sigma}(\mathscr{C}))$.

Proof. Let $c = \sup\{\chi_{st}(G) : G \in \mathsf{Gaifman}_{\Sigma}(\mathscr{C})\} < \infty$. Let $\mathbf{A} \in \mathscr{C}$, let $G = \mathsf{Gaifman}_{\Sigma}(\mathbf{A})$, and let $\gamma : V(G) \to [c]$ be a star coloring of G. In G, any two color classes induce a star forest, which we orient away from their centers. This way we get an orientation \vec{G} of Gsuch that for every vertex v and every in-neighbor u of v, the vertex u is the only neighbor of v with color $\gamma(u)$. Let $R \in \Sigma$ be a relation of arity k. For each $(u_1, \ldots, u_k) \in R(\mathbf{A})$, u_1, \ldots, u_k induce a tournament in \vec{G} . Every tournament has at least one directed Hamiltonian path [Réd34]. We fix one such Hamiltonian path and let $p(u_1, \ldots, u_k)$ be the index of the last vertex in the path. Let $a = p(u_1, \ldots, u_k)$, let $(c_1, \ldots, c_k) = (\gamma(u_1), \ldots, \gamma(u_k))$. Then there exists in G exactly one clique of size k containing u_a with vertices colored c_1, \ldots, c_k , as a consequence of the following claim. (In the claim, $\gamma(K) = \{\gamma(v) : v \in K\}$.)

 \triangleright Claim 3.2. Assume K_1, K_2 are two k-cliques with $\gamma(K_1) = \gamma(K_2)$. If there exists in \vec{G} a directed Hamiltonian path \vec{P} of $\vec{G}[K_1]$ ending at a vertex $v \in K_1 \cap K_2$, then $K_1 = K_2$.

Proof of the claim. We prove the statement by induction on k. If k = 1 the statement is obviously true as $K_1 = K_2 = \{v\}$. Assume that the statement holds for some integer $k \ge 1$, let K_1, K_2 be (k + 1)-cliques with $\gamma(K_1) = \gamma(K_2)$ and assume there exists a directed Hamiltonian path \vec{P} of $\vec{G}[K_1]$ ending at a vertex $v \in K_1 \cap K_2$. Let u be the penultimate vertex of \vec{P} . As $\gamma(K_1) = \gamma(K_2)$, there exists a vertex $u' \in K_2$ with $\gamma(u') = \gamma(u)$. Note that $u' \ne v$ as $\gamma(u) \ne \gamma(v)$. As u is an in-neighbor of v, it is the only neighbor of v in G with color $\gamma(u)$. As K_2 induces a clique, u' is a neighbor of v. Hence, u' = u. Let $K'_1 = K_1 \setminus \{v\}$ and $K'_2 = K_2 \setminus \{v\}$. Then, K'_1 and K'_2 are k-cliques with $\gamma(K'_1) = \gamma(K'_2) = \gamma(K_1) \setminus \{\gamma(v)\}$ and $\vec{P} - \{v\}$ is a directed Hamiltonian path of $\vec{G}[K'_1]$ ending at $u \in K'_1 \cap K'_2$. By the induction hypothesis we have $K'_1 = K'_2$, hence $K_1 = K_2$.

For each relation $R \in \Sigma$ with arity k and each $(u_1, \ldots, u_k) \in R(\mathbf{A})$ we put at $v = u_{p(u_1,\ldots,u_k)}$ a mark $M^R_{\gamma(u_1),\ldots,\gamma(u_k)}$. Then, in the graph G, the vertex v belongs to exactly one clique of size k with vertices colored $\gamma(u_1),\ldots,\gamma(u_k)$, which allows recovering the tuple (u_1,\ldots,u_k) , as all the colors are distinct. We further put at each vertex v a mark $C_{\gamma(v)}$. Then the structure **A** is reconstructed by the transduction $\mathsf{Unfold}_{\Sigma,c}$ defined by the formulas

$$\rho_R(x_1,\ldots,x_k) := \bigvee_{c_1,\ldots,c_k} \Big(\bigwedge_{1 \le j \le k} C_{c_i}(x_i) \wedge \bigwedge_{1 \le i < j \le k} E(x_i,x_j) \wedge \bigvee_{1 \le i \le k} M^R_{c_1,\ldots,c_k}(x_i) \Big). \qquad \Box$$

Note that the condition of Lemma 3.1 is almost tight: if a class \mathscr{C} of undirected graphs contains graphs with arbitrarily large star chromatic number and girth, then the class \mathscr{C} of all orientations of the graphs in \mathscr{C} is not a transduction of \mathscr{C} [NORS20].

Lemma 3.1 will be particularly significant in conjunction with the following results. Recall that a graph G is d-degenerate if every induced subgraph of G contains a vertex of degree at most d, and that a class \mathscr{C} is degenerate if all the graphs in \mathscr{C} are d-degenerate for some d. A class \mathscr{C} of graphs has bounded expansion if, for every integer k, the class of all graphs H whose k-subdivision is a subgraph of some graph in \mathscr{C} is degenerate [NO12].

Theorem 3.3 [BGK⁺21a]. Every degenerate class of graphs with bounded twin-width has bounded expansion.

Theorem 3.4 [NO08]. Every class of graphs with bounded expansion has bounded star chromatic number.

4:11

4. TWIN-MODELS

In this section, we formalize the notions of twin-models and ranked twin-models, which are reminiscent of the "ordered union trees" and "interval biclique partitions" adopted in [BGK⁺21b]. This structure will allow encoding a contraction sequence and to give an alternative definition of twin-width. As mentioned in the introduction, we fix a class \mathscr{C}_0 of binary relational structures with bounded twin-width.

4.1. Twin-models, ranking, layers, and width.

Definition 4.1 (twin-model). Let $\Sigma = (R_1, \ldots, R_k)$ be a binary relational signature. A Σ -twin-model (or simply a twin-model when Σ is clear from the context) is a tuple $(Y, Z_{R_1}, \ldots, Z_{R_k})$ where Y is a rooted binary tree and each Z_{R_i} is a binary relation satisfying the following transversality, minimality, and consistency conditions:

- (transversality) if $(u, v) \in Z_{R_i}$, then u and v are not comparable in the tree order \preceq_Y ;
- (minimality) if $(u, v) \in Z_{R_i}$, then there exists no $(u', v') \neq (u, v)$ with $u' \preceq_Y u, v' \preceq_Y v$ and $(u', v') \in Z_{R_i}$;
- (consistency) if a traversal of a cycle γ in $Y \cup \bigcup_i Z_{R_i}$ respects the *natural orientation* of the Y-edges (that is: the orientation of Y away from the root), then γ contains two consecutive edges in $\bigcup_i Z_{R_i}$.

A twin-model $(Y, Z_{R_1}, \ldots, Z_{R_k})$ defines the Σ -structure \mathbf{A} (or $(Y, Z_{R_1}, \ldots, Z_{R_k})$ is a *twin-model of* \mathbf{A}) if A = L(Y) and, for each $R_i \in \Sigma$, $R_i(\mathbf{A})$ is the set of all pairs (u, v) such that there exists $u' \preceq_Y u$ and $v' \preceq_Y v$ with $(u', v') \in Z_{R_i}$.

Definition 4.2 (ranking, boundaries, layers, and width). Let $(Y, Z_{R_1}, \ldots, Z_{R_k})$ be a twinmodel of a Σ -structure **A** with |A| = n. A ranking τ of the twin-model $(Y, Z_{R_1}, \ldots, Z_{R_k})$ is a mapping from V(Y) to [n] that satisfies the following labeling, monotonicity, and synchronicity conditions:

- (labeling) the function τ restricted to I(Y) is a bijection with [n-1], and is equal to n on L(Y);
- (monotonicity) If $u \prec_Y v$, then $\tau(u) < \tau(v)$;

• (synchronicity) If $(u, v) \in Z_{R_i}$, then $\max(\tau(\pi_Y(u)), \tau(\pi_Y(v))) < \min(\tau(u), \tau(v))$.

A ranked twin-model is a tuple $\mathfrak{T} = (Y, Z_{R_1}, \ldots, Z_{R_k}, \tau)$, where $(Y, Z_{R_1}, \ldots, Z_{R_k})$ is a twin-model, and τ is a ranking of $(Y, Z_{R_1}, \ldots, Z_{R_k})$ (See Figure 5).

For $1 < t \leq n$, the boundary $\partial_t Y$ is the set $\partial_t Y = \{u \in V(Y) \mid \tau(u) \geq t \land \tau(\pi_Y(u)) < t\}$ and the layer \mathbf{L}_t is the Σ^* -structure with vertex set $\partial_t Y$ and relations

$$R_{i}(\mathbf{L}_{t}) = \{(u, v) \in \partial_{t}Y \times \partial_{t}Y \mid \exists u' \preceq_{Y} u, \exists v' \preceq_{Y} v, (u', v') \in Z_{R_{i}}\}$$

$$R_{i}^{*}(\mathbf{L}_{t}) = \{(u, v) \in \partial_{t}Y \times \partial_{t}Y \mid \exists u' \succeq_{Y} u, \exists v' \succeq_{Y} v, (u', v') \neq (u, v)$$
and
$$\{(u', v'), (v', u')\} \cap Z_{R_{i}} \neq \emptyset\}.$$

For t = 1 we define the boundary $\partial_1 Y = \{r(Y)\}$ and the layer \mathbf{L}_1 as the Σ^* -structure with unique vertex r(Y).

The width of the ranked twin-model $\mathfrak{T} = (Y, Z_{R_1}, \dots, Z_{R_k}, \tau)$ is defined as

width(
$$\mathfrak{T}$$
) = max max $\sum_{t \in [n]} \sum_{v \in L_t} |R_i^*(\mathbf{L}_t, v)|,$

Vol. 20:3




Figure 5: A graph G and a ranked twin-model of G. The boundary $\partial_4 Y$ is the set $\{5, g, c, 4\}$ (internal vertices labeled by τ), which can be represented as the set of the yellow zones. The relations of \mathbf{L}_4 are depicted as dotted heavy lines (black for R, red for R^*). The width of this twin-model is 2.

where $R_i^*(\mathbf{L}_t, v)$ denotes the set $\{u : \mathbf{L}_t \models R_i^*(u, v)\}$. Hence, $|R_i^*(\mathbf{L}_t, v)|$ is the degree of v in the symmetric relation R_i^* in \mathbf{L}_t .

At first sight, the consistency condition of a twin-model (of A) may seem contrived. One may for instance wonder if the minimality and consistency conditions are not simply equivalent to the property that every $(u, v) \in R_i(\mathbf{A})$ is realized by a unique unordered pair u', v' with $(u', v') \in Z_{R_i}, u' \preceq_Y u$, and $v' \preceq_Y v$. In case the structure **A** encodes a simple undirected graph G (with signature $\Sigma = (E)$), we would simply impose that the edges of Z_E partition the edges of G into bicliques.

In Figure 6 we give a small example that shows that this property is not strong enough to always yield a ranking. This illustrates why the consistency condition is what we want (no more, no less) and also serves as a visual support for the notions of contraction sequence, twin-model, and ranking.

4.2. From a contraction sequence to a twin-model. In this section, we prove that every d-sequence of a Σ -structure **A** defines a ranked twin-model of **A** with width at most d (See Figure 7).

A d-sequence $\mathbf{A}_n, \ldots, \mathbf{A}_1$ for a Σ -structure \mathbf{A} defines a rooted binary tree Y with vertex set $V(Y) = \bigcup_i A_i$ and set of leaves $L(Y) = A_n$ as follows: for each $i \in [n-1]$ let z_i be the vertex of A_i and u_i, v_i be the vertices of A_{i+1} such that z_i results from the contraction of u_i and v_i in \mathbf{A}_{i+1} . Then $I(Y) = \{z_i : i \in [n-1]\}, r(Y) = z_1$, and the children of z_i in Y are the vertices u_i and v_i .

For each relation $R \in \Sigma$ we define a binary relation Z_R on V(Y) as follows. Let z_i be the vertex of A_i resulting from the contraction of u_i and v_i in A_{i+1} . If $(u_i, v_i) \in R(\mathbf{A}_{i+1})$, then $(u_i, v_i) \in Z_R$. If u_i and v_i are not R-clones for w, then the pairs involving w and u_i or v_i in $R(\mathbf{A}_{i+1})$ are copied in Z_R . Intuitively, Z_R collects the R-relations when they just appear (in the order $\mathbf{A}_1, \ldots, \mathbf{A}_n$). We further define $Z = \bigcup_{R \in \Sigma} Z_R$ and the function $\tau \colon V(Y) \to [n]$ by $\tau(v) = n$ if $v \in L(Y)$ and $\tau(z_i) = i$. Note that for each $i \in [n]$ and non-root vertex v of Y, we have $v \in A_i$ if and only if $\tau(\pi_Y(v)) < i \leq \tau(v)$.

4:12




Figure 6: Left: A 6-vertex graph and a contraction sequence, where the tiny digit in each box indicates the index of contracted vertices when they appear. Center: A twin-model of the graph, where the edges of Z_E are in bold blue, and a ranking (for the internal nodes) of this twin-model that actually matches the contraction sequence. Right: A *flawed twin-model* where the edge set E is indeed partitioned by the pairs of Z_E . Here no ranking is possible: Let α, β, γ be the parents of b, d, f. By synchronicity, and by considering the pairs $(b,\beta), (d,\gamma), \text{ and } (f,\alpha)$, we get that the labeling τ should satisfy $\tau(\alpha) < \tau(\beta), \tau(\beta) < \tau(\gamma), \text{ and } \tau(\gamma) < \tau(\alpha)$, which cannot be realized. There is indeed a cycle $\alpha b\beta d\gamma f$ with all the tree arcs oriented the same way, and without two consecutive edges of Z_E . On the contrary, all such cycles in the central tree have two consecutive edges of Z_E , like 5bd4f has $(b, d), (d, 4) \in Z_E$.




Figure 7: A contraction sequence and the derived ranked tree model.

Lemma 4.3. Every d-sequence $\mathbf{A}_n, \ldots, \mathbf{A}_1$ defines a ranked twin-model with width at most d. *Proof.*

 \triangleright Claim 4.4. The function τ satisfies the labeling, monotonicity, and synchronicity conditions.

Proof of the claim. The first two conditions are straightforward. Let $(u, v) \in Z_R$. Let $i \in [n-1]$ be such that (u, v) appears in \mathbf{A}_i for the first time. As $u, v \in A_i$ we have both $\tau(\pi_Y(u)) < i \leq \tau(u)$ and $\tau(\pi_Y(v)) < i \leq \tau(v)$, i.e., the synchronicity condition holds.

 \triangleright Claim 4.5. The relations Z_R ($R \in \Sigma$) satisfy the minimality and consistency conditions.

Proof of the claim. The minimality condition follows directly from the definition. Let \vec{H} be the oriented graph obtained from Y by orienting all the edges from the root and adding, for each $R \in \Sigma$ and each pair $(u, v) \in Z_R$ the arcs $\pi_Y(u)v$ and $\pi_Y(v)u$ whenever they do not exist. It follows from the monotonicity and synchronicity conditions that \vec{H} is acyclically oriented. Indeed, any arc (x, y) in \vec{H} satisfies $\tau(x) < \tau(y)$. Assume towards a contradiction that in $Y \cup \bigcup_{R \in \Sigma} Z_R$ one can find a cycle γ such that the orientation of the Y-edges is consistent with a traversal of γ and γ does not contain two consecutive edges in $\bigcup_{R \in \Sigma} Z_R$. By replacing in γ each group formed by an edge in $\bigcup_{R \in \Sigma} Z_R$ and its preceding edge in γ (which is in Y) by the corresponding arc in \vec{H} we obtain a circuit in \vec{H} , contradicting its acyclicity. Hence, the relations Z_R satisfy the consistency condition.

From the definition of the width of a ranked twin-model, it is then immediate that the ranked twin-model $(Y, Z_{R_1}, \ldots, Z_{R_k}, \tau)$ derived from a *d*-sequence $\mathbf{A}_n, \ldots, \mathbf{A}_1$ has width at most *d*. This ends the proof of the lemma.

4.3. **Properties of twin-models.** In this section, we establish two properties of twin-models. The first one is the equality of the minimum width of a twin-model with the twin-width of a structure; the second one is that twin-models of structures with bounded twin-width have degenerate Gaifman graphs.

Lemma 4.6. Every twin-model has a ranking, and the twin-width of a Σ -structure **A** is the minimum width of a ranked twin-model of **A**.

Proof. We first prove the first part of the statement.

▷ Claim 4.7. Every twin-model has a ranking.

Proof of the claim. Consider the oriented graph \vec{H} obtained from orienting Y from the root and adding, for each $R \in \Sigma$ and each pair $(u, v) \in Z_R$, an arc $\pi(u)v$ and an arc $\pi(v)u$ (whenever they do not exist). Assume for contradiction that \vec{H} contains a directed cycle. Replace each arc of the form $\pi(u)v$ of this directed cycle (with $(u,v) \in Z_R$) by the path $(\pi(u)u, uv)$ in the twin-model. This way we obtain a closed walk in $Y \cup \bigcup_{R \in \Sigma} Z_R$ traversing all edges of Y away from the root and no two consecutive edges are in $\bigcup_{R \in \Sigma} Z_R$. We show that we can also find a directed cycle in $Y \cup \bigcup_{R \in \Sigma} Z_R$ with this property, contradicting the consistency assumption. Consider a shortest closed walk $W = (e_1, \ldots, e_m)$ with the above property and assume this closed walk is not a directed cycle. Without loss of generality we can assume that (e_1, \ldots, e_k) forms a cycle γ (starting the closed walk at another point if necessary). By minimality of the closed walk, the cycle γ contains two consecutive edges in $\bigcup_{R \in \Sigma} Z_R$. These edges are the edges e_k and e_1 (as otherwise they would be consecutive in W as well). It follows that e_{k+1} does not belong to $\bigcup_{R \in \Sigma} Z_R$ (as it follows e_k in the W). The closed walk $W' = (e_{k+1}, \ldots, e_n)$ does not have two consecutive edges in $\bigcup_{R \in \Sigma} Z_R$ as all the consecutive pairs are consecutive in W, except the pair (e_n, e_{k+1}) (and we know $e_{k+1} \notin \bigcup_{R \in \Sigma} Z_R$). This contradicts the minimality of W. Thus, H is acyclic and a topological ordering of $\overline{H}[I(Y)]$ extends to a labeling $\tau: V(H) \to [n]$ that is bijective between I(Y) and [n-1], equal to n on L(Y), and increasing with respect to every arc of \vec{H} . This directly implies both monotonicity and synchronicity. \triangleleft

The following claim, which asserts that no Z_{R_i} "crosses" the boundaries, will be quite helpful.

 \triangleright Claim 4.8. Let $t \in [n-1]$ and let $u, v \in \partial_t Y$. Then there exists no pair $(u', v') \in Z_{R_i}$ with $u' \prec_Y u$ and $v' \succ_Y v$.

Proof of the claim. Assume $(u', v') \in Z_{R_i}$ and $v' \succ_Y v$. By the synchronicity property we have $\tau(u') > \tau(\pi_Y(v')) \ge \tau(v) \ge t$, contradicting $\tau(u') \le \tau(\pi_Y(u)) < t$.

For a Σ^* -structure **A** and $R \in \Sigma$ we define

$$\overline{R}(\mathbf{A}) = \{(u, v) \in A^2 : \{(u, v), (v, u)\} \cap (R(\mathbf{A}) \cup R^*(\mathbf{A})) \neq \emptyset\}.$$

 \triangleright Claim 4.9. Let $\mathbf{L}_1, \ldots, \mathbf{L}_n$ be the layers of a ranked twin-model \mathfrak{T} of a Σ -structure \mathbf{A} . Then there exists a contraction sequence $\mathbf{A}_n, \ldots, \mathbf{A}_1$ of \mathbf{A} with $A_i = L_i$, and, for each $R \in \Sigma$, $\overline{R}(\mathbf{A}_i) = \overline{R}(\mathbf{L}_i), R(\mathbf{L}_i) \subseteq R(\mathbf{A}_i)$ and $R^*(\mathbf{L}_i) \supseteq R^*(\mathbf{A}_i)$.

Proof of the claim. For $i \in [n-1]$, the Σ -structure \mathbf{A}_i is obtained from \mathbf{A}_{i+1} by contracting the pair of vertices u_i, v_i into w_i , where w_i is the vertex of Y with $\tau(w_i) = i$ and u_i and v_i are the two children of w_i in Y. It is easily checked that $A_i = L_i$. Let z be a vertex of A_i different from w_i . Then $(z, w_i) \in \overline{R}(\mathbf{A}_i)$ if and only if there exists a leaf $w' \succeq_Y w_i$ and a leaf $z' \succeq_Y z$ such that $\{(w', z'), (z', w')\} \cap R(\mathbf{A}) \neq \emptyset$. As \mathfrak{T} is a twin-model of \mathbf{A} this is equivalent to the fact that there exists $w'' \preceq_Y w'$ and $z'' \preceq_Y z'$ with $(w'', z'') \in Z_R$ or $(z'', w'') \in Z_R$. As \preceq_Y is a tree order, w_i and w'' are comparable, as well as z and z''. From this and Claim 4.8 it follows that $(z, w_i) \in \overline{R}(\mathbf{A}_i) \iff \{(z, w_i), (w_i, z)\} \subseteq \overline{R}(\mathbf{L}_i) \iff (z, w_i) \in \overline{R}(\mathbf{L}_i)$, thus $\overline{R}(\mathbf{A}_i) = \overline{R}(\mathbf{L}_i)$.

We now prove $R(\mathbf{L}_i) \subseteq R(\mathbf{A}_i)$ by reverse induction on *i*. For i = n we have $R(\mathbf{L}_i) = R(\mathbf{A}_i) = R(\mathbf{A})$. Let $i \in [n-1]$ and let u_i, v_i, w_i be defined as above. If $(w_i, z) \in R(\mathbf{L}_i)$, then there exists $w' \preceq_Y w_i$ and $z' \preceq_Y z$ with $(w', z') \in Z_R$ thus we have also $(u_i, z) \in R(\mathbf{L}_{i+1})$ and $(v_i, z) \in R(\mathbf{L}_{i+1})$. By induction, we deduce $(u_i, z) \in R(\mathbf{A}_{i+1})$ and $(v_i, z) \in R(\mathbf{A}_{i+1})$. Similarly, if $(z, w_i) \in R(\mathbf{L}_i)$, then $(z, u_i) \in R(\mathbf{A}_{i+1})$ and $(z, v_i) \in R(\mathbf{A}_{i+1})$. Thus, u_i and v_i are *R*-clones for *z*, hence, if $(w_i, z) \in R(\mathbf{L}_i)$, then $(w_i, z) \in R(\mathbf{A}_i)$ and if $(z, w_i) \in R(\mathbf{L}_i)$, then $(z, w_i) \in R(\mathbf{A}_i)$. It follows that we have $R(\mathbf{L}_i) \subseteq R(\mathbf{A}_i)$. Thus, we have

$$R^{*}(\mathbf{L}_{i}) = \overline{R}(\mathbf{L}_{i}) \setminus \left\{ (u, v) : \{(u, v), (v, u)\} \cap R(\mathbf{L}_{i}) = \emptyset \right\}$$
$$\supseteq \overline{R}(\mathbf{A}_{i}) \setminus \left\{ (u, v) : \{(u, v), (v, u)\} \cap R(\mathbf{A}_{i}) = \emptyset \right\}$$
$$= R^{*}(\mathbf{A}_{i}). \qquad \lhd$$

We are now able to complete the proof of the lemma. According to Lemma 4.3, every d-sequence for \mathbf{A} defines a ranked twin-model with width at most d. Conversely, every ranked twin-model for \mathbf{A} with width d' defines a sequence of layers \mathbf{L}_t with $\max_{v \in L_t} \sum_{R_i \in \Sigma} |R_i^*(\mathbf{L}_t, v)| \leq d'$ and, by Claim 4.9, a d'-sequence for \mathbf{A} .

Lemma 4.6 allows introducing the following terminology: the *width* of a twin-model $(Y, Z_{R_1}, \ldots, Z_{R_k})$ is the minimum width of a ranking of $(Y, Z_{R_1}, \ldots, Z_{R_k})$. A twin-model of a Σ -structure **A** is *optimal* if it has the minimum possible width as a twin-model of **A**, which is the twin-width of **A**.

Definition 4.10 (The class \mathscr{T}). The class \mathscr{T} is the class of all optimal twin-models of the Σ -structures in \mathscr{C}_0 .

The following easy remark will be useful.

▷ Claim 4.11. Let $\mathfrak{Y} = (Y, Z_{R_1}, \ldots, Z_{R_k}, \tau)$ be a ranked twin-model of a Σ -structure **A** (with domain A) and let $X \subseteq A$. Let Y' be the subtree of Y induced by all the vertices in X and their pairwise least common ancestors in Y, let Z'_{R_i} be the subset of all pairs in $Z_{R_i} \cap (Y' \times Y')$, and let τ' be the mapping from Y' to [|X|] such that for all $x, y \in V(Y')$ we have $\tau(x) < \tau(y) \iff \tau'(x) < \tau'(y)$. Then $\mathfrak{Y}' = (Y', Z'_{R_1}, \ldots, Z'_{R_k}, \tau')$ is a ranked twin-model of $\mathbf{A}[X]$, whose width is not larger than the one of \mathfrak{Y} .

Lemma 4.12. The Gaifman graph of a ranked twin-model of a Σ -structure with width d is d + 2-degenerate.

Proof. Let $\mathbf{A} = (A, R_1(\mathbf{A}), \dots, R_k(\mathbf{A}))$ be a Σ -structure, let $\mathfrak{T} = (Y, Z_{R_1}, \dots, Z_{R_k}, \tau)$ be a ranked twin-model of \mathbf{A} with width d, and let G be the Gaifman graph of $(Y, Z_{R_1}, \dots, Z_{R_k})$. The ranked twin-model \mathfrak{T} (with layers \mathbf{L}_i) defines a d-sequence $\mathbf{A}_n, \dots, \mathbf{A}_1$, where $A_i = L_i$ (see Lemma 4.6). Let z be the node with $\tau(z) = n - 1$ and let u and v be its children. Each pair in Z_{R_i} containing u (except pairs containing both u and v) gives rise (in \mathbf{A}_{n-1}) to an R_i^* -edge incident to z when contracting u and v. Thus, the degree of u in G is at most d + 2(d for the sum of the degrees in the relations R_i^* , 1 for the pair (u, v) in at least one Z_{R_i} , and 1 for the tree edge (u, z)). Then, in G - u, the vertex v has degree at most d + 1 < d + 2. Now note that by removing u and v from Y, and redefining $\tau(x)$ as $\min(n - 1, \tau(x))$, we get a ranked twin-model of \mathbf{A}_{n-1} (minus R_i^* -edges) with width at most d, whose Gaifman graph is G - u - v. By induction, we deduce that G is d + 2-degenerate.

5. Full twin-models

To reconstruct a Σ -structure **A** from a twin-model $(Y, Z_{R_1}, \ldots, Z_{R_k})$, we make use of the tree order \preceq_Y defined by Y. As this tree order cannot be obtained as a first-order transduction of $(Y, Z_{R_1}, \ldots, Z_{R_k})$ it will be convenient to introduce a variant of twin-models: the *full twinmodel* associated to a twin-model $(Y, Z_{R_1}, \ldots, Z_{R_k})$ is the structure $(V(Y), \prec_Y, Z_{R_1}, \ldots, Z_{R_k})$.

Definition 5.1 (Transduction S and the class \mathscr{F}). The transduction S is the simple interpretation of Σ -structures in full twin-models defined by formulas

$$\rho_0(x) := \neg (\exists y \ y \succ_Y x);$$

$$\rho_{R_i}(x, y) := \exists u \ \exists v \ (u \preceq_Y x) \land (v \preceq_Y y) \land Z_{R_i}(u, v).$$

 \mathscr{F} is the class of all the full twin-models corresponding to the twin-models in \mathscr{T} .

The following lemma follows directly from the definition of a twin-model.

Lemma 5.2. The class \mathscr{C}_0 is a 2-bounded S-transduction of the class \mathscr{F} .

Proof. For all $\mathbf{A} \in \mathscr{F}$, if $\mathbf{T} = (T, \prec, Z_{R_1}, \ldots, Z_{R_k})$ is a full twin-model of \mathbf{A} then $\mathsf{S}(\mathbf{T}) = \mathbf{A}$ and |T| = 2|A| - 1.

Lemma 5.3. Let $\mathfrak{T} = (Y, Z_{R_1}, \ldots, Z_{R_k}, \tau)$ be a ranked twin-model with associated full twinmodel $\mathbf{T} = (V(Y), \prec_Y, Z_{R_1}, \ldots, Z_{R_k})$. Then the twin-width of \mathbf{T} is at most twice the width of the ranked twin-model $(V(Y), \prec_Y, Z_{R_1}, \ldots, Z_{R_k}, \tau)$. *Proof.* Let I_0, I_1 be copies of I(Y) and let $p_i : I(Y) \to I_i$ be the "identity" for i = 0, 1. We define the binary rooted tree \widehat{Y} with vertex set $V(\widehat{Y}) = V(Y) \cup I_1 \cup I_0$, leaf set $L(\widehat{Y}) = V(Y)$, root $r(\widehat{Y}) = p_o(r(Y))$, and parent function

$$\pi_{\widehat{Y}}(x) = \begin{cases} p_1 \circ \pi_Y(x) & \text{if } x \in L(Y) \\ p_0(x) & \text{if } x \in I(Y) \\ p_0 \circ p_1^{-1}(x) & \text{if } x \in I_1 \\ p_1 \circ \pi_Y \circ p_0^{-1}(x) & \text{if } x \in I_0 \setminus \{r(\widehat{Y})\} \\ x & \text{if } x = r(\widehat{Y}) \end{cases}$$

An informal description of \widehat{Y} is that it is obtained by replacing every internal node v of Y by a *cherry* C_v (i.e., a complete binary tree on three vertices) such that one leaf of C_v remains a leaf in \widehat{Y} , the other leaf of C_v is linked to the "children of v", while the root of C_v is linked to the "parent of v" (provided v is not the root of Y).

We further define $\widehat{Z}_{\prec} = \{(v, p_1(v)) : v \in I(Y)\}$ and keep the relations Z_{R_i} as they were defined on \mathfrak{T} . See Figure 8 for an example.




Figure 8: Construction of the twin-model of a twin-model. Each internal vertex of Y defines three vertices of \hat{Y} . Here $I = \{\alpha, \beta, ...\}, I_0 = \{\alpha_0, \beta_0, ...\}$, and $I_1 = \{\alpha_1, \beta_1, ...\}$. Purple (heavy) edges are in Z_R , green (dotted) edges are in \hat{Z}_{\prec} . Note that in the twin-model of the twin-model, only leaves are adjacent by edges in Z_R .

\triangleright Claim 5.4. $\widehat{\mathbf{T}} = (\widehat{Y}, \widehat{Z}_{\prec}, Z_{R_1}, \dots, Z_{R_k})$ is a twin-model of \mathbf{T} .

Proof of the claim. We have $V(Y) = L(\hat{Y})$. The minimality conditions are obviously satisfied for \hat{Z}_{\prec} and Z_{R_i} . Let $\hat{Z} = \hat{Z}_{\prec} \cup \bigcup_i Z_{R_i}$. Consider a cycle $\hat{\gamma}$ in $\hat{Y} \cup \hat{Z}$, with all the edges in \hat{Y} oriented away from the root. Assume for contradiction that no two edges in \hat{Z} are consecutive in $\hat{\gamma}$. Then either $\hat{\gamma}$ contains a directed path of \hat{Y} linking to vertices in L(Y) or a directed path of \hat{Y} linking a vertex in I(Y) to a distinct vertex in V(Y). As no such directed paths exist in \hat{Y} we are led to a contradiction. Hence, $\hat{\mathbf{T}}$ satisfies the consistency condition.

Vol. 20:3

In order to complete our proof, we still need to prove that $\widehat{\mathbf{T}}$ is indeed a twin-model of \mathbf{T} , that is that for every $u, v \in V(Y)$ we have (for all $1 \leq i \leq k$)

$$(u,v) \in Z_{R_i} \text{ (in } \mathbf{T}) \quad \iff \quad \exists u', v' \in V(\widehat{Y}) \quad (u' \preceq_{\widehat{Y}} u) \land (v' \preceq_{\widehat{Y}} v) \land (u',v') \in Z_{R_i} \text{ (in } \widehat{\mathbf{T}})$$
$$u \prec_Y v \text{ (in } \mathbf{T}) \quad \iff \quad \exists u', v' \in V(\widehat{Y}) \quad (u' \preceq_{\widehat{Y}} u) \land (v' \preceq_{\widehat{Y}} v) \land (u',v') \in \widehat{Z}_{\prec} \text{ (in } \widehat{\mathbf{T}})$$

Let $(u, v) \in V(Y)$ and let $1 \le i \le k$..

As in $\widehat{\mathbf{T}}$ the edges in Z_{R_i} link only leaves of \widehat{Y} , we infer that there exists $u' \preceq_{\widehat{Y}} u$ and $v' \preceq_{\widehat{Y}} v$ with $(u', v') \in Z_{R_i}$ (in $\widehat{\mathbf{T}}$) if and only if $(u, v) \in Z_{R_i}$ (in \mathbf{T}); see Figure 8.

As in $\widehat{\mathbf{T}}$ the edges in \widehat{Z}_{\prec} only link some leaf x of \widehat{Y} to $p_1(x)$, we infer that there exists $u' \preceq_{\widehat{Y}} u$ and $v' \preceq_{\widehat{Y}} v$ with $(u', v') \in \widehat{Z}_{\prec}$ (in $\widehat{\mathbf{T}}$) if and only if (up to exchanging u and v) we have $u' = u, v' = p_1(u)$, and $v' \preceq_{\widehat{Y}} v$, that is, if and only if $u \prec_Y v$ (in \mathbf{T}).

Hence, $\widehat{\mathbf{T}}$ is a twin-model of \mathbf{T} .

 \triangleleft

Vol. 20:3

Let n = |L(Y)|. The next claim shows that we have much freedom in defining a ranking for $\widehat{\mathbf{T}}$.

 \triangleright Claim 5.5. If $\hat{\tau} : V(\hat{Y}) \to [2n-1]$ satisfy the labeling and monotonicity conditions, then $\hat{\tau}$ is a ranking of $\hat{\mathbf{T}}$.

Proof of the claim. Assume $(u, v) \in \widehat{Z}_{\prec}$. Then $\pi_{\widehat{Y}}(u) = \pi_{\widehat{Y}}(v)$ hence the synchronicity for \widehat{Z}_{\prec} follows from monotonicity. Assume $(u, v) \in Z_{R_i}$. Then $\widehat{\tau}(u) = \widehat{\tau}(v) = 2n - 1$ hence the synchronicity obviously holds.

We now define $\hat{\tau} : V(\hat{Y}) \to [2n-1]$ as follows: order the vertices $v \in I_1$ by increasing $\tau \circ p_1^{-1}(v)$. For each $v \in I_1$, insert the children of v in I_0 just after v, then add $r(\hat{Y})$ in the very beginning. Numbering the vertices of $I_0 \cup I_1$ according to this order defines $\hat{\tau}$ on $I(\hat{Y})$. We extend this function to the whole $V(\hat{Y})$ by defining $\hat{\tau}(v) = 2n - 1$ for all $v \in L(\hat{Y})$. By construction, the labeling and monotonicity properties hold hence $\hat{\tau}$ is a ranking of $\hat{\mathbf{T}}$.

Consider a time $1 < \hat{t} < 2n - 1$ and let v be the vertex with $\hat{\tau}(v) = \hat{t}$.

Assume for contradiction that some edge (x, y) belongs to Z^*_{\prec} . Then, up to exchanging x and y, x is in I (hence a leaf of \widehat{Y}) and p(x) is a descendant of y (i.e. $p(x) \succ_{\widehat{Y}} y$). Thus, $y \preceq_{\widehat{Y}} \pi_{\widehat{Y}}(p(x)) = \pi_{\widehat{Y}}(x)$, contradicting the necessary condition that x and y are non-comparable in $\preceq_{\widehat{Y}}$. Thus, the degree for Z^*_{\prec} is null.

If $v \in I_1$ we define $t = \tau \circ p_1^{-1}(v)$. Then $\partial_t \widehat{Y} = p_1(\partial_t Y)$ and the degree for $Z_{R_i}^*$ in the layer of \widehat{Y} at time \widehat{t} is at most the degree for $Z_{R_i}^*$ in the layer of Y at time t.

If $v \in I_0$ we define $t = \tau \circ \pi_Y \circ p_0^{-1}(v)$. Then $\partial_t \widehat{Y}$ is $p_1(\partial_t Y)$ in which we remove the parent of v and add v and (maybe) the sibling of v. Compared to the layer at time $\widehat{\tau} \circ \pi_1(\pi_Y \circ p_0^{-1}(v))$, the red degree can only increase because some relations $Z_{R_i}^*$ from a vertex u are incident to v and its sibling. It follows that the maximum $Z_{R_i}^*$ is at most doubled. \Box

From what precedes, we deduce the following theorem, which may be of independent interest.

Theorem 5.6. Let Σ be a binary signature. Every binary Σ -structure with twin-width t has a full twin-width model $\mathbf{T} = (V(Y), \prec_Y, Z_{R_1}, \ldots, Z_{R_k})$ with twin-width at most 2t, associated to a ranked twin-width model $\mathfrak{T} = (V(Y), \prec_Y, Z_{R_1}, \ldots, Z_{R_k}, \tau)$ with width t and d+2-degenerate Gaifman graph.

Proof. The existence of a ranked twin-width model \mathfrak{T} with width t follows from Lemma 4.6. According to Lemma 4.12, \mathfrak{T} has a d + 2-degenerate Gaifman graph and, according to Lemma 5.3, the associated full twin-width model \mathbf{T} has twin-width at most 2t.

6. Ordered twin-models

Recall that a *preordering* of the vertices of a rooted tree is the discovery order of the vertices of a (depth-first search) traversal of the tree starting at its root.

Lemma 6.1. Let \mathscr{O} be the class of binary tree orders and let $\mathscr{Y}^{<}$ be the class of rooted binary trees, with vertices ordered by some preordering. Then there exist simple transductions L and O such that (L, O) is a transduction pairing of \mathscr{O} and $\mathscr{Y}^{<}$.

Proof. We define two simple transductions. The first transduction maps binary tree orders \prec into the preorder defined by some traversal of Y.

 L is defined as follows: we consider a mark M on the vertices and define

$$\rho_E(x,y) := \left((x \prec y) \land \forall v \neg ((x \prec v) \land (v \prec y)) \right) \lor \left((y \prec x) \land \forall v \neg ((y \prec v) \land (v \prec x)) \right)$$
$$\rho_<(x,y) := (x \prec y) \lor \neg (y \preceq x) \land \exists u \exists v \exists w \left(\forall z \left((w \prec z) \rightarrow \neg ((z \prec u) \land (z \prec v)) \right) \land (u \preceq x) \land (v \preceq y) \land (w \prec u) \land (w \prec v) \land \rho_E(u,w) \land \rho_E(v,w) \land M(u) \right).$$

Consider a binary tree order $(V(Y), \prec) \in \mathcal{O}$, and let Y be the rooted binary tree defined by \prec . Recall that the *preordering* of Y defined by some plane embedding of Y (that is to an ordering, for each node v, of the children of v) is a linear order on V(Y) such that for every internal node v of Y, one finds in the ordering the vertex v, then the first child of v and its descendants, then the second child of v and its descendants. Let < be the preordering defined by some plane embedding of Y. (Note, that, despite its name, this is a total order.) Let us mark by M all the nodes of Y that are the first child of their parent. The formula ρ_E defines the cover graph of \prec , thus E is the adjacency relation of Y. Let x, y be nodes of Y. If x = y, then $\rho_{\leq}(x, y)$ does not hold. If x and y are comparable in \prec , then $\rho_{\leq}(x, y)$ is equivalent to $x \prec y$. Otherwise, let w be the infimum of x and y, and let u and v be the children of w such that $u \prec x$ and $v \prec y$. Then $\rho_{\leq}(x, y)$ holds if u is the first child of w, that is, if u is marked. Altogether, we have $\rho_{\leq}(x, y)$ if and only if x < y. Hence, $(V(Y), \prec) \in L(Y^{\leq})$, where Y^{\leq} is Y ordered by \leq .

The transduction O is defined as follows:

$$\rho_{\prec}(x,y) := (x < y) \land \forall z \,\forall w \, \left((x < z) \land (z \le y) \land E(z,w) \right) \to (x \le w).$$

Let $Y^{<}$ be a rooted binary tree Y with preorder < and let \prec be the corresponding tree order. If $x \ge y$, then $\rho_{\prec}(x, y)$ does not hold, so we assume x < y. Assume x is an ancestor of y in Y, then all the vertices z between x and y in the preorder are descendants of x thus any neighbor of these are either descendants of x or x itself thus $\rho_{\prec}(x, y)$ holds. Otherwise, let w be the infimum of x and y in Y and let z be the child of w that is an ancestor of y.

Vol. 20:3

Then z is between x and y is the preorder, z is adjacent to w, but w appears before x in the preorder. Thus, $\rho_{\prec}(x, y)$ does not hold. It follows that $\rho_{\prec}(x, y)$ is equivalent to $x \prec y$ thus $Y^{<} \in O(V(Y), \prec)$. Thus, (L, O) is a transduction pairing of \mathscr{O} and $\mathscr{Y}^{<}$.

Definition 6.2 (The classes $\mathscr{O}_0, \mathscr{Y}_0^<$, and $\mathscr{T}^<$). The class \mathscr{O}_0 is the reduct of the class \mathscr{F} , obtained by keeping only the tree order relation; the class $\mathscr{Y}_0^<$ is the class of all rooted binary trees corresponding to the tree orders in \mathscr{O}_0 ordered by some preordering, so that (L,O) is a transduction pairing of \mathscr{O}_0 and $\mathscr{Y}_0^<$.

The class $\mathscr{T}^{<}$ is the class of ordered twin-models obtained from the twin-models in \mathscr{T} by adding a linear order defined by some preordering of the rooted tree of the tree model.

Note that the natural generalization $(\widehat{L}, \widehat{O})$ of (L, O) is a transduction pairing of \mathscr{F} and $\mathscr{T}^{<}$.

Definition 6.3 (The class \mathscr{G}). The class \mathscr{G} is the class of the Gaifman graphs of the structures in \mathscr{T} .

Lemma 6.4. Let $G = \text{Gaifman}_{\Sigma}$. There exists a simple transduction U such that (G, U) is a transduction pairing of \mathscr{T} and \mathscr{G} .

Proof. According to Lemma 5.3 the class \mathscr{F} has bounded twin-width. According to Theorem 2.2, as $\mathscr{T}^{<}$ is an \widehat{L} -transduction of \mathscr{F} it has bounded twin-width. Thus, the class \mathscr{T} , being a reduct of $\mathscr{T}^{<}$, has bounded twin-width. It follows from Lemma 4.12 that the class $G(\mathscr{T})$ is degenerate, hence, according to Theorem 3.3, it has bounded expansion and, according to Theorem 3.4, bounded star chromatic number (at most c). It follows from Lemma 3.1 that, defining $U = Unfold_{\Sigma,c}$, (G, U) is a transduction pairing of \mathscr{T} and $G(\mathscr{T}) = \mathscr{G}$.

Definition 6.5 (The class $\mathscr{G}^{<}$). The class $\mathscr{G}^{<}$ is the class of ordered graphs obtained from the structures in $\mathscr{T}^{<}$ by applying the natural generalization $\widehat{\mathsf{G}}$ of the interpretation $\mathsf{Gaifman}_{\Sigma}$.

Note that, denoting by \widehat{U} the natural generalization of the transduction U, it follows from Lemma 6.4 that $(\widehat{G}, \widehat{U})$ is a transduction pairing of $\mathscr{T}^{<}$ and $\mathscr{G}^{<}$.

7. Permutations and the main result

When we speak about transductions of permutations, we consider the permutations as defined in Section 2.1. Hence, the language used to define the transduction can use the binary relations $<_1, <_2$, as well as equality.

Lemma 7.1. Let $c \in \mathbb{N}$ and let $\mathscr{S}^{<}$ be a class of ordered graphs, and let \mathscr{S} be the reduct of $\mathscr{S}^{<}$ obtained by forgetting the linear order.

Assume that the graphs in \mathscr{S} have star chromatic number at most c. Then, there exist a copying transduction T_1 with $\mathrm{bf}(\mathsf{T}_1) = c + 1$, a simple transduction T_2 , and a class \mathscr{P} of permutations such that $(\mathsf{T}_1, \mathsf{T}_2)$ is a transduction pairing of $\mathscr{S}^{<}$ and \mathscr{P} .

We give here an informal description of the transductions T_1 and T_2 and refer to Figure 9 for an example: The transduction T_1 is used to compute a permutation σ from an ordered graph $G^<$ and works as follows: we first compute a star coloring γ of G with c colors and orient edges so that bicolored stars are oriented from their centers. Then we blow each vertex into $(u, 1), \ldots, (u, c + 1)$. From this we keep only the vertices of the form (v, c + 1) or of the form (v, i) if v has an in-neighbor colored i. The linear order $<_1$ orders pairs (u, i) by first coordinate first (using <) then by increasing i. The linear order $<_2$ is a succession of intervals ending with a vertex of the form (v, c + 1) (these intervals being ordered according to the order on v); the interval ending with (v, c + 1) contains the pairs $(u, \gamma(v))$, for all the out-neighbors u of v, ordered by first coordinate. The transduction T_2 is used to compute an ordered graph $G^<$ from a permutation σ . It works as follows. First, we mark some elements. These elements will correspond to the vertices of $G^<$, and the linear order < will be the restriction of $<_1$ to these elements. Each vertex v of $G^<$ defines a maximal interval A(v)in $<_1$ ending with v and containing no other marked element, and a maximal interval B(v)in $<_2$ ending with v and containing no other marked element. In $G^<$, a vertex u is adjacent to a vertex $v \neq u$ if A(u) intersects B(v) or A(v) intersects B(u).




Figure 9: In the top, the transduction T_1 (we assume c = 5). In the bottom, the transduction T_2 . In gray, the marked elements of σ , which are the vertices of $G^<$. In $G^<$, an edge links 15 (g in the top) and 28 (v_6 in the top) as A(15) intersects B(28). (The linear order < is the left-hand traversal preorder of the rooted tree.)

Proof. Let $G \in \mathscr{S}$, let $\gamma : V(G) \to [c]$ be a star coloring of G, and let \vec{G} be an orientation of G obtained by orienting all bicolored stars from their roots. We mark a vertex $v \in V(G)$ by M_i if $\gamma(v) = i$ and, for $I \subseteq [c]$, by N_I if I is the set of the γ -colors of the in-neighbors of v. Let C_{c+1} be the (c+1)-blowing transduction. The vertices of $C_{c+1}(G)$ are the pairs $(u,i) \in V(G) \times [c+1]$, and there are new predicates P_j (with $j \in [c+1]$), where $P_j(x)$ holds if x is of the form (u, j), for some $u \in V(G)$. We define $\rho(x) := P_{c+1}(x) \vee \bigvee_{I \subseteq [c]} \bigvee_{i \in I} (N_I(x) \wedge P_i(x))$. (Note that if x is (u, j) then $N_I(x)$ is $N_I(u)$.) Hence $W = \rho(C_{c+1}(G))$ is the union of $V(G) \times \{c+1\}$ and the set of all pairs $(u, i) \in V(G) \times [c]$ such that u has an in-neighbor in \vec{G} with color i. We

consider the subgraph H of $C_{c+1}(G)$ induced by W. For $x \in V(H)$ we define $f_{c+1}(x)$ as x if $P_{c+1}(x)$ or as the (only) ~-neighbor y of x with $P_{c+1}(y)$, that is:

$$f_{c+1}(x) = y \quad \iff \quad P_{c+1}(y) \land ((x = y) \lor (x \sim y)).$$

If x is of the form (u, i) then $f_{c+1}(x)$ is (u, c+1). Then, for $i \in I \subseteq [c]$ and whenever $N_I(x)$ holds, we define $f_i(x)$ as the (only) neighbor y of $f_{c+1}(x)$ in P_{c+1} and M_i . Hence,

$$f_i(x) = y \quad \iff \quad E(f_{c+1}(x), y) \wedge P_{c+1}(y) \wedge M_i(y).$$

We now define the linear orders $<_1$ and $<_2$ as follows (See Figure 9 for an illustration.)

$$\begin{aligned} x <_1 y &\iff \begin{cases} \text{either } f_{c+1}(x) < f_{c+1}(y), \\ \text{or } f_{c+1}(x) = f_{c+1}(y), P_i(x), P_j(y), \text{ and } i < j. \end{cases} \\ x <_2 y &\iff P_i(x), P_j(y), \text{ and } \begin{cases} \text{either } f_i(x) < f_j(y), \\ \text{or } f_i(x) = f_j(y) \text{ and } f_{c+1}(x) < f_{c+1}(y). \end{cases} \end{aligned}$$

(Note that, in the last condition, $f_i(x) = f_j(y)$ implies i = j.)

We call the obtained permutation $\sigma(G^{\leq})$. Note that this permutation depends on some arbitrary choices of star coloring. We further define $\mathscr{P} = \{\sigma(G^{<}) : G^{<} \in \mathscr{S}^{<}\}$. The transduction T_1 is defined as the composition of C_{c+1} , the interpretation reducing the domain to the vertices satisfying $\rho(x)$, then the interpretation defining $<_1, <_2$ and forgetting all the other relations. Hence $\sigma(G^{<}) \in \mathsf{T}_1(G^{<})$.

The definition of T_2 is as follows: we consider a predicate M in such a way that the maximum element of $<_1$ is in M. The domain V of $\mathsf{T}_2(\sigma)$ is $M(\sigma)$. The linear order < is the restriction of $<_1$ to V. Then, x is adjacent to y if there exists $z \notin V$ and $(i, j) \in \{(1, 2), (2, 1)\}$ with $z <_i x, z <_i y$, and no vertex in V is between z and x in $<_i$ and no vertex in V between z and y in $<_j$. It is easily checked that for every $G^{<} \in \mathscr{S}^{<}$ we have $G^{<} \in \mathsf{T}_2(\sigma(G))$ (see Figure 9). According to Lemma 2.1 it follows that (T_1, T_2) is a transduction pairing of $(\mathscr{S}^{<}, \mathscr{P})$.

Theorem 7.2. For every class \mathscr{C}_0 of binary structures with twin-width at most t there exists a proper permutation class $\overline{\mathcal{P}}$, an integer k, and a transduction T, such that \mathcal{C}_0 is a k-bounded T-transduction of $\overline{\mathscr{P}}$. Precisely, for every graph $G \in \mathscr{C}_0$ there is a permutation $\sigma \in \overline{\mathscr{P}}$ on at most k|G| elements with $G \in \mathsf{T}(\sigma)$.

Proof. Let \mathscr{C}_0 be a class of binary structures with twin-width at most t. Let \mathscr{T} be a class of twin-models obtained by optimal contraction sequences of graphs in \mathscr{C}_0 , and let \mathscr{F} be the class of the corresponding full twin-models. According to Lemma 5.3 \mathscr{F} has twin-width at most 2t, moreover, applying the transduction L on \prec we transform \mathscr{F} into the class \mathscr{T}^{\leq} . whose reduct is \mathscr{T} (see Lemma 6.1). Let $\mathscr{G}^{<}$ be the class obtained from $\mathscr{T}^{<}$ be taking the Gaifman graphs of the relations distinct from the linear order, and keeping the linear order, and let \mathscr{G} be the reduct of $\mathscr{G}^{<}$ obtained by forgetting the linear order. Thus $\mathscr{G} = \mathsf{Gaifman}(\mathscr{T})$. As the classes $\mathscr{G}^{<}$ and its reduct \mathscr{G} are transductions of the class \mathscr{F} they have bounded twin-width, by Theorem 2.2. Moreover, the class \mathscr{G} is degenerate hence it has bounded expansion and, in particular, bounded star chromatic number. It follows that we have a transduction pairing of $\mathscr{G}^{<}$ and a class \mathscr{P} of permutations. As the class of all finite graphs has unbounded twin-width and is a transduction of the class of all permutations, the class of all permutations has unbounded twin-width. Thus, as \mathscr{P} has bounded twin-width, it

is a proper class of permutations. From the transduction pairing of $\mathscr{T}^{<}$ and $\mathscr{G}^{<}$ and the one of \mathscr{F} and $\mathscr{T}^{<}$ we deduce that there is a transduction pairing of \mathscr{F} and \mathscr{P} . As \mathscr{C}_0 is a transduction of \mathscr{F} we conclude that \mathscr{C}_0 is a transduction of \mathscr{P} .

Note that the class \mathscr{C}_0 is obviously also a T-transduction of the permutation class $\overline{\mathscr{P}}$ obtained by closing \mathscr{P} under sub-permutations.

Corollary 7.3. Every class of graphs with bounded twin-width contains at most c^n nonisomorphic graphs on n vertices (for some constant c depending on the class).

Proof. Let \mathscr{C}_0 be a class with bounded twin-width. As twin-width is monotone with respect to induced subgraph inclusion, we may assume that \mathscr{C}_0 is hereditary. According to Theorem 7.2, there exists a proper permutation class $\overline{\mathscr{P}}$, an integer k, and a transduction T , such that for every $G \in \mathscr{C}_0$ there is a permutation $\sigma \in \overline{\mathscr{P}}$ on k|G| elements with $G \in \mathsf{T}(G)$. Let m be the number of unary predicates used by the transduction. According to the Marcus-Tardos theorem [MT04], for every proper permutation \mathscr{P} there exists a constant a such that \mathscr{P} contains at most a^n permutations on n elements. For each permutation on n elements, there are 2^{mn} possible choices for the interpretation of the m predicates (as each predicate defines a subset of elements). It follows that \mathscr{C}_0 contains at most $\sum_{i=1}^{kn} a^i 2^{mi} = O((a^k 2^{mk})^n)$ non-isomorphic graphs with at most n vertices. Thus, there exists a constant c such that \mathscr{C}_0 contains at most c^n non-isomorphic graphs with n vertices.

8. Further remarks

It was proved by Bonnet *et al.* [BGO⁺24] that a class of graphs \mathscr{C} has bounded twin-width if and only if it is the reduct of a monadically dependent class of ordered graphs. This implies the following duality type statement for every class $\mathscr{C}^{<}$ of ordered graphs:

$$\exists \text{ permutation } \sigma \text{ with } \operatorname{Av}(\sigma) \longrightarrow \mathscr{C}^{<} \\ \iff \mathscr{C}^{<} - / \gg \mathscr{U} ,$$

where $\operatorname{Av}(\sigma)$ denotes the class of all permutations avoiding the pattern σ , \mathscr{U} denotes the class of all graphs, \longrightarrow means the existence of a transduction, and $-/\gg$ means the non-existence of a transduction.

Interestingly, transductions relate some classical classes of graphs to well studied permutation classes.

Proposition 8.1. A class *C* of graphs has

- bounded linear clique-width if and only if it is a transduction of the class Av(21) of identities;
- bounded clique-width if and only if it is a transduction of the class Av(231) of stack-sortable permutations (or, equivalently, if and only if it is a transduction of the class Av(2413, 3142) of separable permutations).

Proof. It is proved in [Col07] that a class of graphs \mathscr{C} has bounded linear clique-width if and only if it is a transduction of the class of linear orders, that is: if and only if it is a transduction of the class of identities.

It is also proved in [Col07] that a class of graphs \mathscr{C} has bounded clique-width if and only if it is a transduction of the class of (binary) tree orders. The first equivalence of the second item follows from the next claim.

 \triangleright Claim 8.2. Binary tree-orders are transduction equivalent to stack-sortable (i.e. 231-avoiding) permutations.

Proof of the claim. From binary tree-order to 231-avoiding permutations. We let $<_1$ to be the lexicographic order (using two marks) and we define $<_2$ as follows $x <_2 y$ if either x is on the left side at $x \wedge y$ or x < y and the successor x' of x with $x' \leq y$ is on the left side, or x > y and the successor y' of y with $y' \leq x$ is on the right side.

Conversely, from a 231-avoiding permutation $(X, <_1, <_2)$ we define the tree order by $u \wedge y$ is the $<_1$ -maximum element x such that $x <_1 u, x <_1 v$, and x is between u and v in $<_2$.

Let us sketch the second equivalence (with separable permutations): One direction is obvious, as stack-sortable permutations are separable. For the other direction, it is easily shown that separable permutations are transductions of the tree order defined by their separation tree.

Proposition 8.1 shows that two permutations classes (like stack-sortable and separable permutations) may be transduction equivalent but not Wilf-equivalent. To the opposite, it is not difficult to prove that, though they are Wilf-equivalent, the class of 321-avoiding permutations is not a transduction of the class of stack-sortable avoiding permutations. (This follows from the fact that the class of grids is a transduction of Av(321) but has unbounded clique-width.)

The connection between classes of ordered graphs and permutation classes might well be even deeper than what is proved in this paper.

Conjecture 8.3. Every hereditary class $\mathscr{C}^{<}$ of ordered graphs is transduction equivalent to a permutation class.

This conjecture is known to hold if the class $\mathscr{C}^{<}$ is not monadically dependent, as it is then transduction equivalent to the class of all permutations [BGO⁺24]; it also holds if the reduct \mathscr{C} of $\mathscr{C}^{<}$ is biclique-free, as either $\mathscr{C}^{<}$ is not monadically dependent (previous item), or it has bounded twin-width [BGO⁺24]. Then, since biclique-free classes of bounded twinwidth have bounded expansion [BGK⁺21a], and according to Theorem 3.4 and Lemma 7.1 the class $\mathscr{C}^{<}$ is transduction equivalent to a permutation class; finally, it also holds if the reduct \mathscr{C} of $\mathscr{C}^{<}$ is a transduction of a class with bounded expansion as \mathscr{C} is then transduction equivalent to a bounded expansion class \mathscr{D} [GKN⁺20] and this transduction equivalence can be extended to a transduction equivalence of $\mathscr{C}^{<}$ and an expansion $\mathscr{D}^{<}$ of \mathscr{D} , which falls in the previous case.

Acknowledgments

The authors wish to express their gratitude to the referees of a first version of this paper, for their careful reading and their most valuable suggestions.

References

[ABF20] Michael Albert, Mathilde Bouvel, and Valentin Féray. Two first-order logics of permutations. Journal of Combinatorial Theory, Series A, 171:105158, 2020. doi:10.1016/j.jcta.2019.105158.

[BGK⁺21a] Édouard Bonnet, Colin Geniet, Eun Jung Kim, Stéphan Thomassé, and Rémi Watrigant. Twinwidth II: small classes. In Proceedings of the 2021 ACM-SIAM Symposium on Discrete Algorithms (SODA), pages 1977–1996, 2021. doi:10.1137/1.9781611976465.118. Vol. 20:3

- [BGK⁺21b] Édouard Bonnet, Colin Geniet, Eun Jung Kim, Stéphan Thomassé, and Rémi Watrigant. Twinwidth III: Max Independent Set, Min Dominating Set, and Coloring. In Nikhil Bansal, Emanuela Merelli, and James Worrell, editors, 48th International Colloquium on Automata, Languages, and Programming, ICALP 2021, July 12-16, 2021, Glasgow, Scotland (Virtual Conference), volume 198 of LIPIcs, pages 35:1–35:20. Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2021. doi:10.4230/LIPIcs.ICALP.2021.35.
- [BGO⁺24] Édouard Bonnet, Ugo Giocanti, Patrice Ossona de Mendez, Pierre Simon, Stéphan Thomassé, and Szymon Toruńczyk. Twin-width IV: ordered graphs and matrices. Journal of the ACM, 2024. Just Accepted. doi:10.1145/3651151.
- [BKTW22] Édouard Bonnet, Eun Jung Kim, Stéphan Thomassé, and Rémi Watrigant. Twin-width I: tractable FO model checking. J. ACM, 69(1):3:1–3:46, 2022. doi:10.1145/3486655.
- [BNW10] Olivier Bernardi, Marc Noy, and Dominic Welsh. Growth constants of minor-closed classes of graphs. Journal of Combinatorial Theory, Series B, 100(5):468-484, 2010. doi:10.1016/j.jctb. 2010.03.001.
- [Bón12] Miklós Bóna. Combinatorics of permutations. CRC Press, 2012.
- [Cam02] Peter J. Cameron. Homogeneous permutations. The Electronic Journal of Combinatorics, 9(2):R2, 2002. doi:10.37236/1674.
- [Col07] Thomas Colcombet. A combinatorial theorem for trees. In Lars Arge, Christian Cachin, Tomasz Jurdzinski, and Andrzej Tarlecki, editors, Automata, Languages and Programming, 34th International Colloquium, ICALP 2007, Wroclaw, Poland, July 9-13, 2007, Proceedings, volume 4596 of Lecture Notes in Computer Science, pages 901–912. Springer, 2007. doi:10.1007/978-3-540-73420-8_77.
- [GKN⁺20] Jakub Gajarský, Stephan Kreutzer, Jaroslav Nešetřil, Patrice Ossona de Mendez, Michaľ Pilipczuk, Sebastian Siebertz, and Szymon Toruńczyk. First-order interpretations of bounded expansion classes. ACM Transactions on Computational Logic, 21(4):Article 29, 2020. doi: 10.1145/3382093.
- [GM14] Sylvain Guillemot and Dániel Marx. Finding small patterns in permutations in linear time. In Proceedings of the Twenty-Fifth Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2014, Portland, Oregon, USA, January 5-7, 2014, pages 82–101, 2014. doi:10.1137/1. 9781611973402.7.
- [H⁺97] Wilfrid Hodges et al. A shorter model theory. Cambridge university press, 1997.
- [MT04] Adam Marcus and Gábor Tardos. Excluded permutation matrices and the Stanley-Wilf conjecture. J. Comb. Theory, Ser. A, 107(1):153–160, 2004. doi:10.1016/j.jcta.2004.04.002.
- [NO08] Jaroslav Nešetřil and Patrice Ossona de Mendez. Grad and classes with bounded expansion I. Decompositions. European Journal of Combinatorics, 29(3):760-776, 2008. doi:10.1016/j.ejc. 2006.07.013.
- [NO12] Jaroslav Nešetřil and Patrice Ossona de Mendez. Sparsity (Graphs, Structures, and Algorithms), volume 28 of Algorithms and Combinatorics. Springer, 2012. 465 pages.
- [NORS20] Jaroslav Nešetřil, Patrice Ossona de Mendez, Roman Rabinovich, and Sebastian Siebertz. Linear rankwidth meets stability. In Shuchi Chawla, editor, Proceedings of the 2020 ACM-SIAM Symposium on Discrete Algorithms, pages 1180–1199, 2020. doi:10.1137/1.9781611975994.72.
 [Péd24] Légrié Dédzi. Fin herrebingteringhen Setz. Acta Litt. Sugged 7(20,42):07, 1024.
- [Réd34] László Rédei. Ein kombinatorischer Satz. Acta Litt. Szeged, 7(39-43):97, 1934.