

SOME REMARKS ON CONNECTORS AND GROUPOIDS IN GOURSAT CATEGORIES

MARINO GRAN ^a, IDRIS TCHOFFO NGUEFEU ^a, AND DIANA RODELO ^b

^a Institut de Recherche en Mathématique et Physique, Université Catholique de Louvain, Belgium
e-mail address: {marino.gran, idriss.tchoffo}@uclouvain.be

^b Departamento de Matemática, Universidade do Algarve, 3001-501 Coimbra, Portugal
e-mail address: drodelo@ualg.pt

Dedicated to Jiří Adámek on the occasion of his seventieth birthday

ABSTRACT. We prove that connectors are stable under quotients in any (regular) Goursat category. As a consequence, the category $\text{Conn}(\mathbb{C})$ of connectors in \mathbb{C} is a Goursat category whenever \mathbb{C} is. This implies that Goursat categories can be characterised in terms of a simple property of internal groupoids.

Over the last twenty years the property of n -permutability of congruences in a variety of universal algebras has been investigated from a categorical perspective (see [7, 13, 19], for instance, and references therein). When \mathbb{C} is a regular category, the 2-permutability property, usually called the *Mal'tsev* property, is a concept giving rise to a beautiful theory, whose main features are collected in [1]. Many important results still hold when a regular category \mathbb{C} satisfies the strictly weaker property of 3-permutability, namely the *Goursat* property. A nice feature of a (regular) Goursat category \mathbb{C} is that the lattice of equivalence relations on any object in \mathbb{C} is a modular lattice [7], a property that plays a crucial role in commutator theory [10, 17].

The aim of this paper is twofold: first of all we establish some basic properties of Goursat categories in terms of connectors [4], as it was done in [4] for the case of Mal'tsev categories. These results have turned out to be useful to develop a monoidal approach to internal structures [12]. We then give a new characterisation of Goursat categories in terms of properties of (internal) groupoids, on the model of what was done in [11] in the case of Mal'tsev categories.

In the first section, we recall the main properties of Goursat categories that will be needed throughout the paper. In Section 2 we prove that for any Goursat category \mathbb{C} , the category $\text{Equiv}(\mathbb{C})$ of equivalence relations in \mathbb{C} is also a Goursat category (Proposition 2.2, see also [2]). We use this result to give some properties of Goursat categories in terms

Key words and phrases: Goursat categories, 3-permutable varieties, Shifting Lemma, connectors, groupoids.

The third author acknowledges partial financial assistance by Centro de Matemática da Universidade de Coimbra—UID/MAT/00324/2013, funded by the Portuguese Government through FCT/MCTES and co-funded by the European Regional Development Fund through the Partnership Agreement PT2020.

of connectors in Section 3. More precisely, we show that, when \mathbb{C} is a Goursat category, then connectors are stable under quotients in \mathbb{C} (Proposition 3.8), and this implies that the category $\text{Conn}(\mathbb{C})$ of connectors in \mathbb{C} is again a Goursat category (Theorem 3.10).

We conclude the paper by giving a new characterisation of Goursat categories in terms of properties of groupoids and internal categories (Theorem 3.11). It turns out that a regular category \mathbb{C} is a Goursat category if and only if the category $\text{Grpd}(\mathbb{C})$ of groupoids (equivalently, the category $\text{Cat}(\mathbb{C})$ of internal categories) in \mathbb{C} is closed under quotients in the category $\text{RG}(\mathbb{C})$ of reflexive graphs in \mathbb{C} .

1. PRELIMINARIES

In this section we recall some basic definitions and properties of (regular) Goursat categories, needed throughout the article. We shall always assume that the category \mathbb{C} in which we are working is a **regular category**: this means that \mathbb{C} is finitely complete, regular epimorphisms are stable under pullbacks, and kernel pairs have coequalisers. Equivalently, any arrow $f : A \rightarrow B$ has a unique factorisation $f = i \circ r$ (up to isomorphism), where r is a regular epimorphism and i is a monomorphism and this factorisation is pullback stable; the subobject corresponding to i is called the **image** of f .

A **relation** R from X to Y is a subobject $\langle r_1, r_2 \rangle : R \rightarrow X \times Y$. The opposite relation of R , denoted R^o , is the relation from Y to X given by the subobject $\langle r_2, r_1 \rangle : R \rightarrow Y \times X$. A relation R from X to X is called a relation on X . We shall identify a morphism $f : X \rightarrow Y$ with the relation $\langle 1_X, f \rangle : X \rightarrow X \times Y$ and write f^o for its opposite relation. Given another relation $\langle s_1, s_2 \rangle : S \rightarrow Y \times Z$ from Y to Z , one can define the composite relation SR of R and S as the image of the arrow $(r_1 \circ p_1, s_2 \circ p_2) : R \times_Y S \rightarrow X \times Z$, where $(R \times_Y S, p_1, p_2)$ is the pullback of $r_2 : R \rightarrow Y$ along $s_1 : S \rightarrow Y$. With the above notations, any relation $\langle r_1, r_2 \rangle : R \rightarrow X \times Y$ can be seen as the relational composite $r_2 r_1^o$.

The following properties are well known and easy to prove. We collect them in the following lemma:

Lemma 1.1. *Let $f : X \rightarrow Y$ be an arrow in a regular category \mathbb{C} , and let $i \circ r$ be its (regular epimorphism, monomorphism) factorisation. Then:*

- (1) $f^o f$ is the kernel pair of f , thus $1_X \leq f^o f$; moreover, $1_X = f^o f$ if and only if f is a monomorphism;
- (2) $f f^o$ is (i, i) , thus $f f^o \leq 1_Y$; moreover, $f f^o = 1_Y$ if and only if f is a regular epimorphism;
- (3) $f f^o f = f$ and $f^o f f^o = f^o$.

Definition 1.2. A relation (R, r_1, r_2) on an object X is said to be :

- **reflexive** when there is an arrow $r : X \rightarrow R$ such that $r_1 \circ r = 1_X = r_2 \circ r$;
- **symmetric** when there is an arrow $\sigma : R \rightarrow R$ such that $r_2 = r_1 \circ \sigma$ and $r_1 = r_2 \circ \sigma$;
- **transitive** when, by considering the following pullback

$$\begin{array}{ccc} R \times_X R & \xrightarrow{p_2} & R \\ p_1 \downarrow & \lrcorner & \downarrow r_1 \\ R & \xrightarrow{r_2} & X, \end{array}$$

there is an arrow $t : R \times_X R \rightarrow R$ such that $r_1 \circ t = r_1 \circ p_1$ and $r_2 \circ t = r_2 \circ p_2$.

- an **equivalence relation** if R is reflexive, symmetric and transitive.

In particular, a kernel pair $\langle f_1, f_2 \rangle : \text{Eq}(f) \rightrightarrows X \times X$ of a morphism $f : X \rightarrow Y$ is an equivalence relation. The equivalence relations that occur as kernel pairs of some morphism in \mathbb{C} are called **effective**. Let $\text{Equiv}(\mathbb{C})$ be the category whose objects are equivalence relations in \mathbb{C} and arrows from $\langle r_1, r_2 \rangle : R \rightrightarrows X \times X$ to $\langle s_1, s_2 \rangle : S \rightrightarrows Y \times Y$ are pairs (f, g) of arrows in \mathbb{C} making the following diagram commute

$$\begin{array}{ccc} R & \xrightarrow{g} & S \\ r_1 \downarrow & & \downarrow s_1 \\ & & \downarrow s_2 \\ X & \xrightarrow{f} & Y. \end{array}$$

When \mathbb{C} is a regular category, (R, r_1, r_2) is an equivalence relation on X and $f : X \rightarrow Y$ a regular epimorphism, we define the **regular image of R along f** to be the relation $f(R)$ on Y induced by the (regular epimorphism, monomorphism) factorisation $\langle s_1, s_2 \rangle \circ \psi$ of the composite $(f \times f) \circ \langle r_1, r_2 \rangle$:

$$\begin{array}{ccc} R & \xrightarrow{\psi} & f(R) \\ \langle r_1, r_2 \rangle \downarrow & & \downarrow \langle s_1, s_2 \rangle \\ X \times X & \xrightarrow{f \times f} & Y \times Y. \end{array}$$

Note that the regular image $f(R)$ can be obtained as the relational composite $f(R) = fRf^\circ = fr_2r_1^\circ f^\circ$. When R is an equivalence relation, $f(R)$ is also reflexive and symmetric. In a general regular category $f(R)$ is not necessarily an equivalence relation. This is the case in a *Goursat category* (Theorem 1.4).

Definition 1.3 [8, 7]. A regular category \mathbb{C} is called a **Goursat category** when the equivalence relations in \mathbb{C} are 3-permutable, i.e. $RSR = SRS$ for any pair of equivalence relations R and S on the same object.

The following characterisation will be useful in the sequel:

Theorem 1.4 [7]. *A regular category \mathbb{C} is a Goursat category if and only if for any regular epimorphism $f : X \rightarrow Y$ and any equivalence relation R on X , the regular image $f(R) = fRf^\circ$ of R along f is an equivalence relation.*

There are many important algebraic examples of Goursat categories. Indeed, by a classical theorem in [18], a variety of universal algebras is a Goursat category precisely when its theory has two ternary operations r and s such that the identities $r(x, y, y) = x$, $r(x, x, y) = s(x, y, y)$ and $s(x, x, y) = y$ hold. Accordingly, the categories of groups, abelian groups, modules over some fixed ring, crossed modules, quasi-groups, rings, associative algebras, Heyting algebras and implication algebras are all Goursat categories.

Any regular *Mal'tsev* category is a Goursat category, thus, in particular, so is any semi-abelian category.

Many interesting properties of Goursat categories can be found in the literature (see [7, 15, 16] and references therein). In particular, the following characterisations will be useful for the development of this work:

Consequently, a morphism (f, g) is a monomorphism in $\text{Equiv}(\mathbb{C})$ if and only if f and g are monomorphisms in \mathbb{C} . When \mathbb{C} is a Goursat category, a similar property holds with respect to regular epimorphisms:

Lemma 2.1. *Let R and S be two equivalence relations in a Goursat category \mathbb{C} and $(f, g) : R \rightarrow S$ a morphism*

$$\begin{array}{ccc} R & \xrightarrow{g} & S \\ r_1 \downarrow & & \downarrow s_1 \\ X & \xrightarrow{f} & Y \\ r_2 \downarrow & & \downarrow s_2 \end{array} \quad (2.2)$$

in $\text{Equiv}(\mathbb{C})$. Then (f, g) is a regular epimorphism in $\text{Equiv}(\mathbb{C})$ if and only if f and g are regular epimorphisms in \mathbb{C} .

Proof. When f and g are regular epimorphisms in \mathbb{C} , it is not difficult to check that (f, g) is necessarily the coequaliser of its kernel pair in $\text{Equiv}(\mathbb{C})$ given in (2.1) (one uses the fact that $g = \text{coeq}(g_1, g_2)$ and $f = \text{coeq}(f_1, f_2)$ in \mathbb{C}).

Conversely, let (f, g) be a morphism in $\text{Equiv}(\mathbb{C})$ as in (2.2) that is a regular epimorphism in $\text{Equiv}(\mathbb{C})$. Consider the kernel pairs of f and g , the (regular epimorphism, monomorphism) factorisation $f = i \circ q$ of f , and the regular image $(q(R), t_1, t_2)$ of (R, r_1, r_2) along q . We obtain the following commutative diagram

$$\begin{array}{ccccc} \text{Eq}(g) & \xrightarrow{g_1} & R & \xrightarrow{g} & S \\ & \xrightarrow{g_2} & \downarrow r_1 & \searrow \alpha & \downarrow s_1 \\ & & R & & q(R) \\ & & \downarrow r_2 & & \downarrow t_1 \\ \text{Eq}(f) & \xrightarrow{f_1} & X & \xrightarrow{f} & Y \\ & \xrightarrow{f_2} & \downarrow q & \searrow i & \downarrow t_2 \\ & & X & & Z \\ & & & & \downarrow i \end{array} \quad (2.3)$$

where $(q(R), t_1, t_2) \in \text{Equiv}(\mathbb{C})$ (by Theorem 1.4) and (i, j) is the morphism in $\text{Equiv}(\mathbb{C})$ such that $(i, j) \circ (q, \alpha) = (f, g)$. Note that j is induced from the fact that $(i \times i) \circ \langle t_1, t_2 \rangle \circ \alpha$ is the (regular epimorphism, monomorphism) factorisation of $\langle s_1, s_2 \rangle \circ g$, thus it is a monomorphism

$$\begin{array}{ccc} R & \xrightarrow{\alpha} & q(R) \\ g \downarrow & \searrow j & \downarrow (i \times i) \circ \langle t_1, t_2 \rangle \\ S & \xrightarrow{\langle s_1, s_2 \rangle} & Y \times Y \end{array}$$

From the fact that (f, g) is the coequaliser of its kernel pair in $\text{Equiv}(\mathbb{C})$ it easily follows that (i, j) is an isomorphism in $\text{Equiv}(\mathbb{C})$. This implies that f and g are regular epimorphisms in \mathbb{C} . \square

Proposition 2.2. *Equiv(\mathbb{C}) is a Goursat category whenever \mathbb{C} is.*

Proof. As mentioned above, the category $\text{Equiv}(\mathbb{C})$ is finitely complete because \mathbb{C} is so. Lemma 2.1 implies that regular epimorphisms in $\text{Equiv}(\mathbb{C})$ are stable under pullbacks since regular epimorphisms are stable in \mathbb{C} , and regular epimorphisms in $\text{Equiv}(\mathbb{C})$ are “levelwise” regular epimorphisms. The existence of the (regular epimorphism, monomorphism) factorisation of a morphism (f, g) as in (2.2) in the category $\text{Equiv}(\mathbb{C})$ follows from the construction of diagram (2.3): the (regular epimorphism, monomorphism) factorisation $f = i \circ q$ of f in \mathbb{C} gives rise to the (regular epimorphism, monomorphism) factorisation $g = j \circ \alpha$ of g in \mathbb{C} . Thus $(q, \alpha) \circ (i, j)$ is the (regular epimorphism, monomorphism) factorisation of (f, g) in $\text{Equiv}(\mathbb{C})$. To see that $\text{Equiv}(\mathbb{C})$ has the Goursat property one uses Theorem 1.4: to check that the regular image of an equivalence relation in the category $\text{Equiv}(\mathbb{C})$ is again an equivalence in $\text{Equiv}(\mathbb{C})$ one mainly uses the same (“levelwise”) property in the category \mathbb{C} . \square

3. CONNECTORS AND GROUPOIDS IN GOURSAT CATEGORIES

In this section we prove that connectors are stable under quotients in any Goursat category \mathbb{C} . We then define the category $\text{Conn}(\mathbb{C})$ of connectors in \mathbb{C} whose objects are pairs of equivalence relations equipped with a connector, and prove that $\text{Conn}(\mathbb{C})$ is a Goursat category whenever the base category \mathbb{C} is. We conclude by giving a new characterisation of Goursat categories in terms of properties of groupoids and internal categories.

Definition 3.1. Let (R, r_1, r_2) and (S, s_1, s_2) be two equivalence relations on an object X and $R \times_X S$ the pullback of r_2 along s_1 . A **connector** [4] between R and S is an arrow $p: R \times_X S \rightarrow X$ in \mathbb{C} such that

- (1) $xSp(x, y, z)Rz$;
- (2) $p(x, x, y) = y$;
- (3) $p(x, y, y) = x$;
- (4) $p(x, y, p(z, u, v)) = p(p(x, y, z), u, v)$,

when each term is defined.

Remark 3.2. Given two regular epimorphisms $d: X \twoheadrightarrow Y$ and $c: X \twoheadrightarrow Z$, a connector on the effective equivalence relations $\text{Eq}(d)$ and $\text{Eq}(c)$ is the same thing as an internal *pregroupoid* in the sense of Kock [20, 21] (see also the introduction of [5], for instance, for a comparison between these two related notions and some additional references). In the context of Mal’tsev or Goursat categories connectors are useful to develop a centrality theory of non-effective equivalence relations.

Example 3.3. *If ∇_X is the largest equivalence relation on an object X , then an associative Mal’tsev operation*

$$p: X \times X \times X \rightarrow X$$

is precisely a connector between ∇_X and ∇_X .

Connectors provide a way to distinguish groupoids amongst reflexive graphs:

Proposition 3.4 [9]. *Given a reflexive graph*

$$X_1 \begin{array}{c} \xrightarrow{d} \\ \xleftarrow{e} \\ \xrightarrow{c} \end{array} X_0$$

in a finitely complete category \mathcal{C} (i.e. $d \circ e = 1_{X_0} = c \circ e$) then the connectors between $\text{Eq}(c)$ and $\text{Eq}(d)$ are in bijections with the groupoid structures on this reflexive graph.

It is well known that Goursat categories satisfy the so-called *Shifting Property* [17, 6]. In this context connectors are unique when they exist (Theorem 2.13 and Proposition 5.1 in [6]): accordingly, for a given pair of equivalence relations on the same object the fact of having a connector becomes a *property*.

Definition 3.5. Let R and S be two equivalence relations on an object X . A **double equivalence relation** on R and S is given by an object $C \in \mathbb{C}$ equipped with two equivalence relations $(\pi_1, \pi_2) : C \rightrightarrows R$ and $(p_1, p_2) : C \rightrightarrows S$ such that the following diagram

$$\begin{array}{ccc} C & \begin{array}{c} \xrightarrow{\pi_1} \\ \xrightarrow{\pi_2} \end{array} & R \\ p_1 \downarrow & \begin{array}{cc} p_2 & r_1 \end{array} & \downarrow r_2 \\ S & \begin{array}{c} \xrightarrow{s_1} \\ \xrightarrow{s_2} \end{array} & X \end{array}$$

commutes (in the “obvious” way).

A double equivalence relation C on R and S is called a **centralizing relation** [9] when the square

$$\begin{array}{ccc} C & \xrightarrow{\pi_1} & R \\ p_1 \downarrow & \lrcorner & \downarrow r_1 \\ S & \xrightarrow{s_1} & X \end{array}$$

is a pullback. Under this assumption it follows that any of the commutative squares in the definition of a centralizing relation is a pullback.

The following lemma gives the relationship between connectors and centralizing relations.

Lemma 3.6 [4]. *If \mathbb{C} is a category with finite limits and R and S are two equivalence relations on the same object X , then the following conditions are equivalent:*

- (i) *there exists a connector between R and S ;*
- (ii) *there exists a centralizing relation on R and S .*

When \mathbb{C} is a Mal'tsev category, R and S are equivalence relations on an object X with a connector and $i : I \rightarrow X$ is a monomorphism, then the inverse images $i^{-1}(R)$ and $i^{-1}(S)$ also have a connector [4]. We establish a similar property for Goursat categories, with respect to regular epimorphisms:

Proposition 3.7. *Let \mathbb{C} be a Goursat category, R and S two equivalence relations on an object X , and let $f : X \rightarrow Y$ be a regular epimorphism. If there exists a connector between R and S , then there exists a connector between the regular images $f(R)$ and $f(S)$.*

Proof. Suppose that there exists a connector between R and S . This implies that there exists a centralizing relation $(C, (\pi_1, \pi_2), (p_1, p_2))$ on R and S . Consider the regular image

$(f(R), a, b)$ and $(f(S), c, d)$ of R and S along f . We obtain the following diagram

$$\begin{array}{ccccc}
 C & \xrightarrow{\alpha} & f_R(C) & & \\
 \downarrow p_1 & \searrow \pi_2 & \downarrow \beta_2 & & \\
 & \downarrow \pi_1 & \downarrow \beta_1 & & \\
 R & \xrightarrow{f_R} & f(R) & & \\
 \downarrow p_2 & \searrow r_1 & \downarrow a & & \\
 & \downarrow r_2 & \downarrow b & & \\
 S & \xrightarrow{f_S} & f(S) & & \\
 \downarrow s_1 & \searrow s_2 & \downarrow c & & \\
 X & \xrightarrow{f} & Y & &
 \end{array}
 \tag{3.1}$$

where $(f_R(C), \beta_1, \beta_2)$ is the regular image of the equivalence relation (C, π_1, π_2) along the regular epimorphism f_R . The fact that the square

$$\begin{array}{ccc}
 C & \xrightarrow{\alpha} & f_R(C) \\
 f_S p_1 \downarrow & \swarrow \alpha_1 & \downarrow \langle a\beta_1, a\beta_2 \rangle \\
 f(S) & \xrightarrow{\langle c, d \rangle} & Y \times Y
 \end{array}$$

commutes, α is a strong epimorphism and $\langle c, d \rangle$ is a monomorphism, implies the existence of an arrow $\alpha_1 : f_R(C) \rightarrow f(S)$ making the above diagram commute. Similarly, from the commutativity of the third diagram

$$\begin{array}{ccc}
 C & \xrightarrow{\alpha} & f_R(C) \\
 f_S p_2 \downarrow & \swarrow \alpha_2 & \downarrow \langle b\beta_1, b\beta_2 \rangle \\
 f(S) & \xrightarrow{\langle c, d \rangle} & Y \times Y
 \end{array}$$

we obtain an arrow $\alpha_2 : f_R(C) \rightarrow f(S)$.

The relations $(f_R(C), \beta_1, \beta_2)$, $(f(R), a, b)$ and $(f(S), c, d)$ are all equivalence relations by Theorem 1.4. It is then easy to check that the relation $(f_R(C), \alpha_1, \alpha_2)$ is an equivalence relation on $f(S)$. In fact, the morphism $\langle \alpha_1, \alpha_2 \rangle : f_R(C) \rightarrow f(S) \times f(S)$ is a monomorphism since $\langle c \times c, d \times d \rangle \circ \langle \alpha_1, \alpha_2 \rangle = \langle a, b \rangle \times \langle a, b \rangle \circ \langle \beta_1, \beta_2 \rangle$. So, $\langle \alpha_1, \alpha_2 \rangle$ is the regular image of $\langle p_1, p_2 \rangle$ along f_S , thus it is an equivalence relation on $f(S)$ by Theorem 1.4.

By assumption all the left squares of (3.1) are pullbacks, so it follows that all the right squares of (3.1) are pullbacks as well by Theorem 1.6 (ii). Then $(f_R(C), (\alpha_1, \alpha_2), (\beta_1, \beta_2))$ is a centralizing relation on $f(R)$ and $f(S)$. By Lemma 3.6 there is a connector between $f(R)$ and $f(S)$. \square

We are now going to show that the category whose objects are pairs of equivalence relations equipped with a connector is a Goursat category whenever the base category is a Goursat category. For this, let us first fix some notation: if \mathbb{C} is a finitely complete category, we write $2\text{-Eq}(\mathbb{C})$ for the category whose objects (R, S, X) are pairs of equivalence relations

R and S on the same object X

$$R \begin{array}{c} \xrightarrow{r_1} \\ \xrightarrow{r_2} \end{array} X \begin{array}{c} \xleftarrow{s_1} \\ \xleftarrow{s_2} \end{array} S$$

and arrows are triples (f_R, f_S, f) making the following diagram commute:

$$\begin{array}{ccc} R & \begin{array}{c} \xrightarrow{r_1} \\ \xrightarrow{r_2} \end{array} & X \begin{array}{c} \xleftarrow{s_1} \\ \xleftarrow{s_2} \end{array} & S \\ f_R \downarrow & & f \downarrow & \downarrow f_S \\ \bar{R} & \begin{array}{c} \xrightarrow{\bar{r}_1} \\ \xrightarrow{\bar{r}_2} \end{array} & \bar{X} \begin{array}{c} \xleftarrow{\bar{s}_1} \\ \xleftarrow{\bar{s}_2} \end{array} & \bar{S}. \end{array} \quad (3.2)$$

We write $\text{Conn}(\mathbb{C})$ for the category whose objects (R, S, X, p) are pairs of equivalence relations R and S on an object X with a given connector $p : R \times_X S \rightarrow X$; arrows in $\text{Conn}(\mathbb{C})$ are arrows in $2\text{-Eq}(\mathbb{C})$ respecting the connectors. This means that, given a diagram (3.2) where both (R, S, X) and $(\bar{R}, \bar{S}, \bar{X})$ are in $\text{Conn}(\mathbb{C})$, with $p : R \times_X S \rightarrow X$ and $\bar{p} : \bar{R} \times_{\bar{X}} \bar{S} \rightarrow \bar{X}$ the corresponding connectors, then the diagram

$$\begin{array}{ccc} R \times_X S & \xrightarrow{\bar{f}} & \bar{R} \times_{\bar{X}} \bar{S} \\ p \downarrow & & \downarrow \bar{p} \\ X & \xrightarrow{f} & \bar{X} \end{array}$$

commutes, where \bar{f} is the natural map induced by the universal property of the pullback $\bar{R} \times_{\bar{X}} \bar{S}$.

We say that a subcategory \mathbb{P} is *closed under (regular) quotients* in a category \mathbb{Q} if, for any regular epimorphism $f : A \rightarrow B$ in \mathbb{Q} such that $A \in \mathbb{P}$, then $B \in \mathbb{P}$.

Proposition 3.8. *If \mathbb{C} is a Goursat category, then $\text{Conn}(\mathbb{C})$ is a full subcategory of $2\text{-Eq}(\mathbb{C})$, that is closed in $2\text{-Eq}(\mathbb{C})$ under quotients.*

Proof. The fullness of the forgetful functor $\text{Conn}(\mathbb{C}) \rightarrow 2\text{-Eq}(\mathbb{C})$ follows from Corollary 5.2 in [6], by taking into account the fact that any Goursat category satisfies the Shifting Property.

Let us then consider a regular epimorphism in $2\text{-Eq}(\mathbb{C})$

$$\begin{array}{ccc} R & \begin{array}{c} \xrightarrow{r_1} \\ \xrightarrow{r_2} \end{array} & X \begin{array}{c} \xleftarrow{s_1} \\ \xleftarrow{s_2} \end{array} & S \\ f_R \downarrow & & f \downarrow & \downarrow f_S \\ \bar{R} & \begin{array}{c} \xrightarrow{\bar{r}_1} \\ \xrightarrow{\bar{r}_2} \end{array} & \bar{X} \begin{array}{c} \xleftarrow{\bar{s}_1} \\ \xleftarrow{\bar{s}_2} \end{array} & \bar{S} \end{array}$$

(this means that f , f_R and f_S are regular epimorphisms in \mathbb{C}) such that its domain (R, S, X) belongs to $\text{Conn}(\mathbb{C})$. The equalities $f(R) = \bar{R}$ and $f(S) = \bar{S}$, together with Proposition 3.7, imply that there exists a connector between \bar{R} and \bar{S} . \square

Lemma 3.9. *Let \mathbb{D} be a finitely complete category, and \mathbb{C} a full subcategory of \mathbb{D} closed in \mathbb{D} under finite limits and quotients. Then:*

- (1) \mathbb{C} is regular whenever \mathbb{D} is regular.
- (2) \mathbb{D} is a Goursat category whenever \mathbb{C} is a Goursat category.

Proof. The (regular epimorphism, monomorphism) factorisation in \mathbb{D} of an arrow in \mathbb{C} is also its factorisation in \mathbb{C} , since \mathbb{C} is closed in \mathbb{D} under quotients. Since finite limits in \mathbb{C} are calculated as in \mathbb{D} , it follows that regular epimorphisms are stable under pullbacks. Now the second statement easily follows from the fact that the composition of relations is computed in the same way in \mathbb{C} and in \mathbb{D} . \square

Theorem 3.10. *If \mathbb{C} is a Goursat category then $\text{Conn}(\mathbb{C})$ is a Goursat category.*

Proof. Using similar arguments as those given in the proof of Proposition 2.2 with respect to $\text{Equiv}(\mathbb{C})$, one may deduce that $2\text{-Eq}(\mathbb{C})$ is a Goursat category. The result then follows from Proposition 3.8 and Lemma 3.9. \square

We finally prove that internal categories and groupoids can be used to characterise Goursat categories. Recall that an **internal category** in a category \mathbb{C} with pullbacks is a reflexive graph with a multiplication $m: X_1 \times_{X_0} X_1 \rightarrow X_1$

$$X_1 \times_{X_0} X_1 \begin{array}{c} \xrightarrow{p_1} \\ \xrightarrow{m} \\ \xrightarrow{p_2} \end{array} X_1 \begin{array}{c} \xrightarrow{d} \\ \xleftarrow{e} \\ \xrightarrow{c} \end{array} X_0,$$

(where $X_1 \times_{X_0} X_1$ is the pullback of d and c) such that:

- $d \circ m = d \circ p_2$, $c \circ m = c \circ p_1$, $m \circ \langle e \circ d, 1_{X_1} \rangle = 1_{X_1} = m \circ \langle 1_{X_1}, e \circ c \rangle$;
- $m \circ (1 \times m) = m \circ (m \times 1)$.

An internal category

$$X_1 \times_{X_0} X_1 \begin{array}{c} \xrightarrow{p_1} \\ \xrightarrow{m} \\ \xrightarrow{p_2} \end{array} X_1 \begin{array}{c} \xrightarrow{d} \\ \xleftarrow{e} \\ \xrightarrow{c} \end{array} X_0,$$

is a **groupoid** when there is an additional morphism $i: X_1 \rightarrow X_1$ satisfying the axioms:

- $d \circ i = c$, $c \circ i = d$;
- $m \circ \langle i, 1_{X_1} \rangle = e \circ c$ and $m \circ \langle 1_{X_1}, i \rangle = e \circ d$.

We write $\text{Cat}(\mathbb{C})$ for the category of internal categories in \mathbb{C} (and internal functors as morphisms), $\text{Grpd}(\mathbb{C})$ for the category of groupoids in \mathbb{C} , and $\text{RG}(\mathbb{C})$ for the category of reflexive graphs in \mathbb{C} (with obvious morphisms).

An equivalence relation is a special kind of groupoid, where its domain and codomain morphisms are jointly monomorphic; also any reflexive and transitive relation is in particular an internal category. If \mathbb{C} is a Goursat category, then any reflexive and transitive relation is an equivalence relation or, equivalently, any internal category is a groupoid (Theorem 1 in [22]). Then Theorem 1.4, which could equivalently be stated through the property that $\text{Equiv}(\mathbb{C})$ (or the category of reflexive and transitive relations in \mathbb{C}) is closed in the category of reflexive relations in \mathbb{C} under quotients, has an *extended* counterpart given below. This characterisation leads to the observation that the structural aspects of Goursat categories mainly concern groupoids (rather than equivalence relations).

Theorem 3.11. *Let \mathbb{C} be a regular category. Then the following conditions are equivalent:*

- (i) \mathbb{C} is a Goursat category;
- (ii) $\text{Grpd}(\mathbb{C})$ is closed in $\text{RG}(\mathbb{C})$ under quotients;
- (iii) $\text{Cat}(\mathbb{C})$ is closed in $\text{RG}(\mathbb{C})$ under quotients.

Proof.

(i) \Rightarrow (ii) Let

$$\begin{array}{ccc} X_1 & \xrightarrow{g} & X'_1 \\ d \downarrow \uparrow c & & d' \downarrow \uparrow c' \\ X_0 & \xrightarrow{f} & X'_0 \end{array}$$

be a regular epimorphism (f, g) in $\text{RG}(\mathbb{C})$ (which means that f and g are regular epimorphisms in \mathbb{C}), with

$$X_1 \begin{array}{c} \xrightarrow{d} \\ \xleftarrow{c} \\ \xrightarrow{c} \end{array} X_0$$

a groupoid in \mathbb{C} . By Proposition 3.4, there exists a connector between $\text{Eq}(d)$ and $\text{Eq}(c)$. Let $\text{Eq}(d)$, $\text{Eq}(c)$, $\text{Eq}(d')$ and $\text{Eq}(c')$ be the kernel pairs of the arrows d , c , d' and c' , respectively. Let $\lambda : \text{Eq}(d) \rightarrow \text{Eq}(d')$ and $\beta : \text{Eq}(c) \rightarrow \text{Eq}(c')$ be the arrows induced by the universal property of kernel pairs $\text{Eq}(d')$ and $\text{Eq}(c')$, respectively. By Theorem 1.5, λ and β are regular epimorphisms, so that $g(\text{Eq}(d)) = \text{Eq}(d')$ and $g(\text{Eq}(c)) = \text{Eq}(c')$. By Proposition 3.7 there is then a connector between $\text{Eq}(d')$ and $\text{Eq}(c')$, thus

$$X'_1 \begin{array}{c} \xrightarrow{d'} \\ \xleftarrow{c'} \\ \xrightarrow{c'} \end{array} X'_0$$

is a groupoid (Proposition 3.4).

- (ii) \Rightarrow (i) This implication follows from Theorem 1.4 and the fact that equivalence relations are in particular groupoids (whose domain and codomain morphisms are jointly monomorphic).
- (i) \Rightarrow (iii) This implication follows from (i) \Rightarrow (ii) and the fact that $\text{Grpd}(\mathbb{C}) \cong \text{Cat}(\mathbb{C})$ in a Goursat context, as recalled above.
- (iii) \Rightarrow (i) Let (R, r_1, r_2) be an equivalence relation on X , $f : X \twoheadrightarrow Y$ a regular epimorphism and $(f(R), t_1, t_2)$ the regular image of R along f

$$\begin{array}{ccc} R & \xrightarrow{g} & f(R) \\ r_1 \downarrow \downarrow r_2 & & t_1 \downarrow \downarrow t_2 \\ X & \xrightarrow{f} & Y \end{array}$$

$(f(R), t_1, t_2)$ is reflexive and symmetric being the image of the equivalence relation R along a regular epimorphism f . By assumption, $(f(R), t_1, t_2)$ is an internal category, thus it is an equivalence relation. It follows that \mathbb{C} is a Goursat category (by Theorem 1.4). \square

Remark 3.12. Observe that Theorem 3.11 also implies that $\text{Grpd}(\mathbb{C})$ and $\text{Cat}(\mathbb{C})$ are Goursat categories whenever \mathbb{C} is, again thanks to Lemma 3.9, the category $\text{RG}(\mathbb{C})$ obviously being a Goursat category. This simplifies and slightly extends Proposition 4.3 in [14], where the existence of coequalizers in \mathbb{C} was assumed.

Remark 3.13. A result analogous to Theorem 3.11 holds in the Mal'tsev context: a category \mathbb{C} is a Mal'tsev category if and only if $\text{Grpd}(\mathbb{C})$ (or, equivalently, $\text{Cat}(\mathbb{C})$) is closed in $\text{RG}(\mathbb{C})$ under subobjects [3]. Together with the comments made before Proposition 3.7 we observe the existence of a sort of duality between Mal'tsev categories and Goursat categories.

Similar results hold for Mal'tsev categories with respect to monomorphisms and for Goursat categories with respect to regular epimorphisms.

REFERENCES

- [1] F. Borceux and D. Bourn, *Mal'cev, protomodular, homological and semi-abelian categories*, Mathematics and its Applications, 566. Kluwer Academic Publishers, Dordrecht, 2004. xiv+479 pp. ISBN: 1-4020-1961-0
- [2] D. Bourn, *Internal equivalence relations, modular formula and Goursat condition in non-regular context*, Technical Report 526 of the L.M.P.A. (2016).
- [3] D. Bourn, *Mal'cev categories and fibration of pointed objects*, Appl. Categ. Structures 4 (1996) 307327.
- [4] D. Bourn and M. Gran, *Centrality and connectors in Maltsev categories*, Algebra Universalis 48 (2002), no. 3, 309-331.
- [5] D. Bourn and M. Gran, *Centrality and normality in protomodular categories*, CT2000 Conference (Como). Theory Appl. Categ. 9 (2001/02), 151-165.
- [6] D. Bourn and M. Gran, *Categorical aspects of modularity*. Galois theory, Hopf algebras, and semiabelian categories, 77100, Fields Inst. Commun., 43, Amer. Math. Soc., Providence, RI, 2004.
- [7] A. Carboni, G.M. Kelly, M.C. Pedicchio, *Some remarks on Mal'tsev and Goursat categories*, Appl. Categ. Structures 1 (1993), no. 4, 385-421.
- [8] A. Carboni, J. Lambek, M.C. Pedicchio, *Diagram chasing in Mal'cev categories*, J. Pure Appl. Algebra 69 (1991) 271-284.
- [9] A. Carboni, M.C. Pedicchio, N. Pirovano, *Internal graphs and internal groupoids in Mal'cev categories*, Category theory 1991 (Montreal, PQ, 1991), 97-109, CMS Conf. Proc., 13, Amer. Math. Soc., Providence, RI, 1992.
- [10] R. Freese and R. McKenzie, *Commutator theory for congruence modular varieties*, Cambridge University Press, Cambridge (1987).
- [11] M. Gran, *Internals categories in Mal'cev categories*, Special volume on the occasion of the 60th birthday of Professor Michael Barr (Montreal, QC, 1997). J. Pure Appl. Algebra 143 (1999), no. 1-3, 221-229.
- [12] M. Gran, C. Heunen, and S. Tull, *Monoidal methods for groupoids and connectors in categorical algebra*, in preparation.
- [13] M. Gran, M. C. Pedicchio *n-permutables locally finitely presentable categories*, Theory Appl. Categ. 8 (2001), 1-15.
- [14] M. Gran and D. Rodelo, *A universal construction in Goursat categories*, Cah. Topol. Géom. Différ. Catég. 49 (2008), no. 3, 196-208.
- [15] M. Gran and D. Rodelo, *A new characterisation of Goursat categories*, Appl. Categ. Structures 20 (2012), no. 3, 229-238.
- [16] M. Gran and D. Rodelo, *Beck-Chevalley condition and Goursat categories*, J. Pure Appl. Algebra, 221 (2017) 2445-2457.
- [17] H.P. Gumm, *Geometrical methods in congruence modular algebras*, Mem. Amer. Math. Soc. 45 (1983), no. 286, viii+79 pp.
- [18] J. Hagemann and A. Mitschke, *On n-permutable congruences*, Algebra Universalis 3 (1973), 8-12.
- [19] Z. Janelidze, D. Rodelo and T. Van der Linden, *Hagemann's theorem for regular categories*, J. Homotopy Relat. Struct. 9(1) (2014) 55-66.
- [20] A. Kock, *Generalized fibre bundles*, Lecture Notes in Mathematics 1348, Springer-Verlag (1988) 194-207.
- [21] A. Kock, *Fibre bundles in general categories*, J. Pure Appl. Algebra, 56 (1989) 233-245.
- [22] N. Martins-Ferreira, D. Rodelo and T. Van der Linden, *An observation on n-permutability*, Bull. Belg. Math. Soc. Simon Stevin 21 (2014), no. 2, 223-230.