

EVERY METRIC SPACE IS SEPARABLE IN FUNCTION REALIZABILITY

ANDREJ BAUER AND ANDREW SWAN

Andrej Bauer, Faculty of Mathematics and Physics, University of Ljubljana, Jadranska 19, 1000
Ljubljana, Slovenia
e-mail address: Andrej.Bauer@andrej.com

Andrew Swan, Institute for Logic, Language and Computation, University of Amsterdam, Science
Park 107, 1098 XG Amsterdam, Netherlands
e-mail address: wakelin.swan@gmail.com

ABSTRACT. We first show that in the function realizability topos $\text{RT}(\mathcal{K}_2)$ every metric space is separable, and every object with decidable equality is countable. More generally, working with synthetic topology, every T_0 -space is separable and every discrete space is countable. It follows that intuitionistic logic does not show the existence of a non-separable metric space, or an uncountable set with decidable equality, even if we assume principles that are validated by function realizability, such as Dependent and Function choice, Markov's principle, and Brouwer's continuity and fan principles.

Are there any uncountable sets with decidable equality in constructive mathematics, or at least non-separable metric spaces? We put these questions to rest by showing that in the function realizability topos all metric spaces are separable, and consequently all sets with decidable equality countable. Therefore, intuitionistic logic does not show existence of non-separable metric spaces, even if we assume principles that are validated by function realizability, among which are the Dependent and Function choice, Markov's principle, and Brouwer's continuity and fan principles.

1. FUNCTION REALIZABILITY

We shall work with the realizability topos $\text{RT}(\mathcal{K}_2)$, see [8, §4.3], which is based on Kleene's function realizability [3]. We carry out the bulk of the argument in the internal language of the topos, which is intuitionistic logic with several extra principles, cf. Proposition 1.1.

We write \mathbf{N} and \mathbf{R} for the objects of the natural numbers and the real numbers, respectively. The Baire space is the object $\mathbf{B} = \mathbf{N}^{\mathbf{N}}$ of infinite number sequences. It is metrized by the metric $u : \mathbf{B} \times \mathbf{B} \rightarrow \mathbf{R}$ defined by

$$u(\alpha, \beta) = \lim_{n \rightarrow \infty} 2^{-\min\{k \leq n \mid \alpha_k \neq \beta_k\}}.$$

Key words and phrases: Constructive mathematics, function realizability, metric spaces, synthetic topology.

The first author acknowledges that this material is based upon work supported by the Air Force Office of Scientific Research under award number FA9550-17-1-0326.

If the first index at which α and β differ is k , then $u(\alpha, \beta) = 2^{-k}$.

Proposition 1.1. *The realizability topos $\text{RT}(\mathcal{K}_2)$ validates the following principles:*

- (1) Countable choice: *a total relation on \mathbb{N} has a choice map.*
- (2) Extended function choice: *if $S \subseteq \mathbb{B}$ is $\neg\neg$ -stable then every total relation on S has a choice map.*
- (3) Extended continuity principle: *if $S \subseteq \mathbb{B}$ is $\neg\neg$ -stable then every map $S \rightarrow \mathbb{B}$ is continuous.*
- (4) Excluded middle for predicates on \mathbb{N} : *if $\phi(n)$ is a formula whose only parameter is $n \in \mathbb{N}$, then $\forall n \in \mathbb{N}. \phi(n) \vee \neg\phi(n)$.*

Proof.

- (1) Every realizability topos validates Countable choice, and [8, Prop. 4.3.2] does so specifically for $\text{RT}(\mathcal{K}_2)$.
- (2) Recall that a subobject $S \subseteq \mathbb{B}$ is $\neg\neg$ -stable when $\neg\neg(\alpha \in S)$ implies $\alpha \in S$ for all $\alpha \in \mathbb{B}$. The realizability relation on such an S is inherited by that of \mathbb{B} , i.e., the elements of S are realized by Kleene's associates. The argument proceeds the same way as [8, Prop. 4.3.2], which shows that choice holds in the case $S = \mathbb{B}$.
- (3) Once again, if $S \subseteq \mathbb{B}$ is $\neg\neg$ -stable, then maps $S \rightarrow \mathbb{B}$ are realized by Kleene's associates, just like maps from $\mathbb{B} \rightarrow \mathbb{B}$. The argument proceeds the same way as continuity of maps $\mathbb{B} \rightarrow \mathbb{B}$ in [8, Prop. 4.3.4].
- (4) Let us first show that, for a formula $\psi(n)$ whose only parameter is $n \in \mathbb{N}$, the sentence

$$\forall n \in \mathbb{N}. \neg\neg\psi(n) \Rightarrow \psi(n) \tag{1.1}$$

is realized. The formula $\psi(n)$ is interpreted as a subobject of \mathbb{N} , which is represented by a map $f : \mathbb{N} \rightarrow \mathcal{P}(\mathbb{N}^{\mathbb{N}})$. Using a bit of (external) classical logic and Countable choice we obtain a map $c : \mathbb{N} \rightarrow \mathbb{N}^{\mathbb{N}}$ such that

$$\forall n \in \mathbb{N}. f(n) \neq \emptyset \Rightarrow c(n) \in f(n),$$

which says that c can be used to build a realizer for (1.1).

Now, given a formula $\phi(n)$ whose only parameter is $n \in \mathbb{N}$, take $\psi(n)$ to be $\phi(n) \vee \neg\phi(n)$. Because $\neg\neg(\phi(n) \vee \neg\phi(n))$ holds, (1.1) reduces to the desired statement $\forall n \in \mathbb{N}. \phi(n) \vee \neg\phi(n)$. \square

Note that the last part of the previous proposition does *not* state the validity of the internal statement

$$\forall \phi \in \Omega^{\mathbb{N}}. \forall n \in \mathbb{N}. \phi(n) \vee \neg\phi(n).$$

Indeed, such a statement cannot be valid in any realizability topos because it implies excluded middle (given $p \in \Omega$, consider $\phi(n) \equiv p$). Rather, we have a *schema* which holds for each formula $\phi(n)$.

2. METRIC SPACES IN FUNCTION REALIZABILITY

Henceforth we argue in the internal language of $\text{RT}(\mathcal{K}_2)$. A minor but common complication arises because the internal language does not allow quantification over all objects of the topos. Thus, when we make a statement $\phi_{(X,d)}$ about all metric spaces (X, d) in the topos, this is to be understood schematically: given any object X and morphism $d : X \times X \rightarrow \mathbb{R}$, if the topos validates “ d is a metric” then it also validates $\phi_{(X,d)}$.

A consequence of Countable choice is that the object of reals \mathbf{R} is a continuous image of \mathbf{B} , for instance by composing any bijection $\mathbf{B} \cong \mathbf{Z}^{\mathbf{N}}$ with the surjection $\mathbf{Z}^{\mathbf{N}} \rightarrow \mathbf{R}$ defined by taking $\alpha \in \mathbf{Z}^{\mathbf{N}}$ to $\alpha_0 + \sum_{k=0}^{\infty} 2^{-k} \cdot \max(-1, \min(1, \alpha_{k+1}))$.

We follow the convention that a set X is *countable* if there exists a surjection $\mathbf{N} \rightarrow X + 1$, and we refer to such a surjection as an *enumeration* of X . Note in particular that the empty set is countable, and that if X is inhabited, then it is countable if and only if there is a surjection $\mathbf{N} \rightarrow X$.

Recall that a metric space (X, d) is *separable* if there exists a countable subset $D \subseteq X$ such that, for all $x \in X$ and $k \in \mathbf{N}$, there exists $y \in D$ such that $d(x, y) \leq 2^{-k}$.

Proposition 2.1. *Suppose (X, d_X) is a separable metric space and $q : X \rightarrow Y$ is a surjection onto a metric space (Y, d_Y) such that $d_Y \circ (q \times q) : X \times X \rightarrow \mathbf{R}$ is continuous with respect to the metric d_X . Then Y is separable.*

Proof. Let $D \subseteq X$ be a countable dense subset of X . We claim that its image $q(D)$ is dense in Y . Consider any $y \in Y$ and $k \in \mathbf{N}$. There is $x \in X$ such that $q(x) = y$. Because $d_Y \circ (q \times q)$ is continuous at (x, x) there exists $m \in \mathbf{N}$ such that, for all $z \in X$, if $d_X(x, z) \leq 2^{-m}$ then

$$d_Y(q(x), q(z)) = |d_Y(q(x), q(z)) - d_Y(q(x), q(x))| \leq 2^{-k}.$$

Since D is dense there exists $z \in D$ such that $d_X(x, z) \leq 2^{-m}$, hence $d_Y(y, q(z)) = d_Y(q(x), q(z)) \leq 2^{-k}$. \square

Proposition 2.2. *For any subobject $S \rightarrow \mathbf{B}$, the topos $\mathbf{RT}(\mathcal{K}_2)$ validates the statement that (S, u) is a separable metric space.*

Proof. Let \mathbf{N}^* be the object of finite sequences of numbers. We write $|a|$ for the length n of a sequence $a = (a_0, \dots, a_{n-1})$. For $\alpha \in \mathbf{B}$ and $k \in \mathbf{N}$, let $\bar{\alpha}(k) = (\alpha_0, \dots, \alpha_{k-1})$ be the prefix of α of length k . Given a finite sequence $a = (a_0, \dots, a_{n-1})$, let $a0^\omega$ be its padding by zeroes,

$$a0^\omega = (a_0, \dots, a_{n-1}, 0, 0, 0, \dots).$$

Notice that $u(\bar{\alpha}(k)0^\omega, \alpha) \leq 2^{-k}$ for all $\alpha \in \mathbf{B}$ and $k \in \mathbf{N}$.

Because \mathbf{N}^* is isomorphic to \mathbf{N} , we may apply Excluded middle for predicates on \mathbf{N} to establish

$$\forall a \in \mathbf{N}^* . (\exists \alpha \in S . u(a0^\omega, \alpha) \leq 2^{-|a|}) \vee \neg(\exists \alpha \in S . u(a0^\omega, \alpha) \leq 2^{-|a|}).$$

By Countable choice there is a map $c : \mathbf{N}^* \rightarrow S + 1$ such that, for all $a \in \mathbf{N}^*$, if there exists $\alpha \in S$ with $u(a0^\omega, \alpha) \leq 2^{-|a|}$ then $c(a) \in S$ and $u(a0^\omega, c(a)) \leq 2^{-|a|}$. We claim that c enumerates a dense sequence in S . To see this, consider any $\alpha \in S$ and $k \in \mathbf{N}$. Because $u(\bar{\alpha}(k+1)0^\omega, \alpha) \leq 2^{-k-1}$, we have $c(\bar{\alpha}(k+1)) \in S$ and therefore

$$\begin{aligned} u(\alpha, c(\bar{\alpha}(k+1))) &\leq \\ u(\alpha, \bar{\alpha}(k+1)0^\omega) + u(\bar{\alpha}(k+1)0^\omega, c(\bar{\alpha}(k+1))) &\leq 2^{-k-1} + 2^{-k-1} = 2^{-k}. \quad \square \end{aligned}$$

Proposition 2.3. *A metric space is separable if its carrier is the quotient of a $\neg\neg$ -stable subobject of \mathbf{B} .*

Proof. Suppose (X, d) is a metric space such that there exist a $\neg\neg$ -stable $S \subseteq \mathbf{B}$ and a surjection $q : S \rightarrow X$. By Proposition 2.2, the space (S, u) is separable. We may apply

Proposition 2.1, provided that $d \circ (q \times q) : S \times S \rightarrow \mathbb{R}$ is continuous with respect to u . By the Extended function choice, $d \circ (q \times q)$ factors through a continuous surjection $\mathbb{B} \twoheadrightarrow \mathbb{R}$ as

$$\begin{array}{ccc} S \times S & \xrightarrow{f} & \mathbb{B} \\ & \searrow & \swarrow \\ & d \circ (q \times q) & \mathbb{R} \end{array}$$

By the Extended Continuity principle the map f is continuous, hence $d \circ (q \times q)$ is continuous, too. \square

Before proceeding we review the notion of a *modest* object [8, Def. 3.2.23]: X is modest when it has $\neg\neg$ -stable equality and is orthogonal to the object $\nabla 2$ [8, Prop. 3.2.22].¹ The modest objects are, up to isomorphism, the quotients by $\neg\neg$ -stable equivalence relations of $\neg\neg$ -stable subobjects of the underlying partial combinatory algebra, which in our case is the Baire space \mathbb{B} . The powers and the subobjects of a modest set are modest (see the remark after [8, Def. 3.2.23] about applicability of [8, Prop. 3.2.19] to modest objects).

Theorem 2.4. *Every metric space in $\text{RT}(\mathcal{K}_2)$ is separable.*

Proof. Consider a metric space (X, d) in $\text{RT}(\mathcal{K}_2)$. The object of reals \mathbb{R} is modest because it has $\neg\neg$ -stable equality and is a quotient of \mathbb{B} . Its power \mathbb{R}^X is modest, and because the transpose of the metric $\tilde{d} : X \rightarrow \mathbb{R}^X$ embeds X into \mathbb{R}^X , the carrier X is modest, therefore a quotient of a $\neg\neg$ -stable subobject of \mathbb{B} . We may apply Proposition 2.3. \square

Theorem 2.5. *In $\text{RT}(\mathcal{K}_2)$ every object with decidable equality is countable.*

Proof. The precise statement is: for any object X in $\text{RT}(\mathcal{K}_2)$, $\text{RT}(\mathcal{K}_2)$ validates the statement “if X has decidable equality then X is countable”.

We argue internally. If equality on X is decidable then we may define the discrete metric $d : X \times X \rightarrow \mathbb{R}$ by

$$d(x, y) = \begin{cases} 0 & \text{if } x = y, \\ 1 & \text{otherwise} \end{cases}$$

Because (X, d) is separable it contains a countable dense subset $D \subseteq X$. But then $D = X$, because for any $x \in X$, there is $y \in D$ such that $d(x, y) < 1/2$, which implies $x = y$. \square

The upshot of Theorems 2.4 and 2.5 is that in constructive mathematics a non-separable metric space cannot be constructed, and neither can an uncountable set with decidable equality, for such constructions could be interpreted in $\text{RT}(\mathcal{K}_2)$ to give counterexamples to the theorems.

The theorems fail for *families* of objects. For instance, while for any specific subobject $S \rightarrow \mathbb{N}$ the statement “ S is countable” is valid, the internal statement “every subobject of \mathbb{N} is countable” is invalid. It is a variation of *Kripke’s schema* [7] which together with Markov’s principle implies Excluded middle, as follows.

Proposition 2.6. *If every subobject of \mathbb{N} is countable and Markov’s principle holds, then Excluded middle holds as well.*

¹The results of [8, §3.2] refer specifically to the effective topos, but are easily adapted to any realizability topos, as long as one replaces \mathbb{N} with the underlying partial combinatory algebra, especially in [8, Def. 3.2.17].

Proof. Let us show that the stated assumptions imply that every truth value p is $\neg\neg$ -stable. From an enumeration of the subobject $\{n \in \mathbb{N} \mid p\}$, which exists by assumption, we may construct $f : \mathbb{N} \rightarrow \{0, 1\}$ such that p is equivalent to $\exists n \in \mathbb{N}. f(n) = 1$. Markov's principle says that such a statement is $\neg\neg$ -stable. \square

Corollary 2.7. *The statement “not every subobject of \mathbb{N} is countable” is valid in $\text{RT}(\mathcal{K}_2)$.*

Proof. A realizability topos validates Markov's principle and the negation of the law of excluded middle. \square

3. INTRINSIC T_0 -SPACES ARE SEPARABLE IN FUNCTION REALIZABILITY

We now generalize our result to *synthetic topology*. We refer the reader to [1, 4, 2] for comprehensive accounts of synthetic topology, and recall only those concepts that are needed for our results. The *Rosolini dominance* [6] is the subobject of Ω , defined by

$$\Sigma = \{p : \Omega \mid \exists \alpha : \{0, 1\}^{\mathbb{N}}. p \Leftrightarrow (\exists n \in \mathbb{N}. \alpha_n = 1)\}.$$

The object Σ is modest and can be thought of as an analogue of the *Sierpiński space*, which classifies open subspaces in the category of topological spaces and continuous maps. This observation is the starting point of synthetic topology, where the exponential Σ^X is taken to be the *intrinsic topology* of an object X . Thus a subobject of X is *intrinsically open* when its characteristic map $X \rightarrow \Omega$ factors through the inclusion $\Sigma \rightarrow \Omega$. We call the elements of Σ the *open truth values*.

With regards to metric spaces, a fundamental question is how the metric and intrinsic topologies relate. Because the relation $<$ on \mathbb{R} is intrinsically open [1, Prop. 2.1], every metric open ball is intrinsically open. We note that the converse holds for $\neg\neg$ -stable subobjects of \mathbb{B} .

Proposition 3.1. *Every intrinsically open subspace of a $\neg\neg$ -stable subobject $S \rightarrow \mathbb{B}$ is a union of open balls.*

Proof. The object Σ is a quotient of \mathbb{B} by the map $\beta \mapsto (\exists n. \beta(n) = 1)$. Consider an intrinsically open subset $U \subseteq S$, with characteristic map $u : S \rightarrow \Sigma$. By Extended function choice u factors through the quotient $\mathbb{B} \rightarrow \Sigma$ as a map $\bar{u} : S \rightarrow \mathbb{B}$. Moreover, by the Extended continuity principle the map \bar{u} is continuous. Suppose $\alpha \in S$ and $u(\alpha) = \top$. There is $n \in \mathbb{N}$ such that $\bar{u}(\alpha)(n) = 1$. By continuity of \bar{u} there is an open ball $B(\alpha, r)$ centered at α such that, for all $\beta \in B(\alpha, r)$, we have $\bar{u}(\beta)(n) = 1$, therefore $\alpha \in B(\alpha, r) \subseteq U$, as required. \square

Many standard topological notions may be formulated in synthetic topology. For instance, an object X is an (*intrinsic*) T_0 -space when the transpose $X \rightarrow \Sigma^{\Sigma^X}$ of the evaluation map $\Sigma^X \times X \rightarrow \Sigma$ is a monomorphism, i.e., when two points in X are equal if they have the same open neighborhoods. Metric spaces are T_0 -spaces because metric open balls are intrinsically open.

In classical topology arbitrary unions of opens are open, but this is not so in synthetic topology. We say that I is *overt* if Σ is closed under I -indexed unions. In logical form overtness says that for every $u : I \rightarrow \Sigma$ the truth value $\exists i \in I. u(i)$ is open. Because Σ is a lattice, Kuratowski-finite objects are overt. The natural numbers \mathbb{N} are overt as well, by an application of Countable choice.

In general overtness transfers along (*intrinsically*) *dense* maps, which are maps $f : X \rightarrow Y$ such that, for all $u : Y \rightarrow \Sigma$,

$$(\exists y \in Y . u(y)) \iff (\exists x \in X . u(f(x))).$$

Clearly, if X is overt and $f : X \rightarrow Y$ is dense then Y is overt.

Say that X is (*intrinsically*) *separable* when there exists a dense map $\mathbb{N} \rightarrow X$. Because \mathbb{N} is overt, all intrinsically separable objects are overt.

Lemma 3.2. *Every $\neg\neg$ -stable subobject of \mathbf{B} is intrinsically separable.*

Proof. By Proposition 3.1, the intrinsic and metric topologies of a $\neg\neg$ -stable subobject $S \rightarrow \mathbf{B}$ agree, hence so do both kinds of separability. In Proposition 2.2 we showed that S is metrically separable. \square

As noted earlier, every metric space is a T_0 -space, and intrinsic separability implies metric separability. Thus, the following is a generalization of Theorem 2.4.

Theorem 3.3. *In $\mathbf{RT}(\mathcal{K}_2)$, every intrinsically T_0 -space is intrinsically separable.*

Proof. Suppose X is a T_0 -space. Then it is a subobject of the modest object Σ^{Σ^X} , and so it is also modest. Hence there exists a surjection $q : S \twoheadrightarrow X$ from a $\neg\neg$ -stable subobject $S \rightarrow \mathbf{B}$. By Lemma 3.2 there is a dense map $f : \mathbb{N} \rightarrow S$. But a dense map followed by a surjection is dense, and so $q \circ f$ witnesses intrinsic separability of X . \square

Finally, a generalization of Theorem 2.5 is readily available. An object X is (*intrinsically*) *discrete* when the equality relation on X is open, or equivalently, when every singleton in X is open.

Theorem 3.4. *In $\mathbf{RT}(\mathcal{K}_2)$, every intrinsically discrete space is countable.*

Proof. An intrinsically discrete space X is a T_0 -space because in X singletons are open. By Theorem 3.3 there is a dense map $f : \mathbb{N} \rightarrow X$, which must be a surjection because, again, singletons are open. \square

4. CONCLUSION

In view of our results, it is natural to wonder what goes wrong intuitionistically with the classical non-separable spaces, such as ℓ^∞ and $L^\infty[0, 1]$. Do they somehow become separable? One of several things can happen. In $\mathbf{RT}(\mathcal{K}_2)$ every map $[0, 1] \rightarrow \mathbf{R}$ is uniformly continuous so that $L^\infty[0, 1]$ is just the space of uniformly continuous maps with the supremum norm, which of course is separable. On the other hand, ℓ^∞ cannot even be constructed, at least not using the classical definition of the norm

$$\|a\|_\infty = \sup_{n \in \mathbb{N}} |a_n|.$$

For a bounded $a : \mathbb{N} \rightarrow \mathbf{R}$ the supremum need not exist, so that further restrictions on a are required. The most generous attempt would collect into ℓ^∞ all the sequences for which $\|a\|_\infty$ exists — but doing so would break the vector space structure, and consequently the definition of the metric. We could also observe that $\|a\|_\infty$ is a well-defined *lower* real number, i.e., we may give it as a lower Dedekind cut. How much of the mathematics of ℓ^∞ , and similar spaces, one can recover this way was studied by Fred Richman [5],

Lastly, we remark that the results are quite closely tied to $\text{RT}(\mathcal{K}_2)$ because of the non-computable nature of the Excluded middle for predicates on \mathbb{N} from Proposition 1.1 (apply the principle to the statement “the n -th Turing machine halts” to obtain the halting oracle). A close cousin of $\text{RT}(\mathcal{K}_2)$ is the Kleene-Vesley topos, which is defined as the relative realizability topos on the partial combinatory subalgebra of \mathcal{K}_2 consisting of the computable functions [8, §4.5]. Because in this topos all statements must be realized by computable maps, the last part of the proof of Proposition 1.1 fails. Indeed, an uncountable object with decidable equality is readily available, just take a subset of \mathbb{N} which is not computably enumerable, and therefore not countable internally to the topos.

REFERENCES

- [1] Andrej Bauer and Davorin Lešnik. Metric spaces in synthetic topology. *Annals of Pure and Applied Logic*, 163(2):87–100, 2012.
- [2] Martín Hötzel Escardó. Synthetic topology of data types and classical spaces. *Electronic Notes in Theoretical Computer Science*, 87:21–156, 2004.
- [3] Stephen Cole Kleene and Richard Eugène Vesley. *The Foundations of Intuitionistic Mathematics, especially in relation to recursive functions*. North-Holland Publishing Company, 1965.
- [4] Davorin Lešnik. *Synthetic Topology and Constructive Metric Spaces*. PhD thesis, University of Ljubljana, 2010.
- [5] Fred Richman. Generalized real numbers in constructive mathematics. *Indagationes Mathematicae*, 9(4):595 – 606, 1998.
- [6] Giuseppe Rosolini. *Continuity and Effectiveness in Topoi*. PhD thesis, University of Oxford, 1986.
- [7] Anne Sjerp Troelstra. *Principles of intuitionism*. Number 95 in Lecture Notes in Mathematics. Berlin, 1969.
- [8] Jaap van Oosten. *Realizability: An Introduction to its Categorical Side*, volume 152 of *Studies in Logic and the Foundations of Mathematics*. Elsevier, 2008.