

CO-C.E. SPHERES AND CELLS IN COMPUTABLE METRIC SPACES

ZVONKO ILJAZOVIĆ

Department of Mathematics, Faculty of Science, University of Zagreb, Bijenička 30, 10 000 Zagreb,
Croatia
e-mail address: zilj@math.hr

ABSTRACT. We investigate conditions under which a co-computably enumerable set in a computable metric space is computable. Using higher-dimensional chains and spherical chains we prove that in each computable metric space which is locally computable each co-computably enumerable sphere is computable and each co-c.e. cell with co-c.e. boundary sphere is computable.

1. INTRODUCTION

A closed subset of \mathbb{R}^m is said to be *computable* if it can be effectively approximated by a finite set of points with rational coordinates with arbitrary given precision on arbitrary given bounded region of \mathbb{R}^m . A closed subset of \mathbb{R}^m is said to be *co-computably enumerable* (*co-c.e.*) if its complement can be effectively covered by open balls. Each computable set is co-c.e. On the other hand, there exist co-c.e. sets which are not computable. In fact, while each nonempty computable set contains computable points, there exists a nonempty co-c.e. set which contains no computable points ([9]). Although the implication

$$S \text{ co-computably enumerable} \Rightarrow S \text{ computable} \quad (1.1)$$

does not hold in general, there are certain conditions under which it does hold. The following result has been proved in [7]:

- (i) if $S \subseteq \mathbb{R}^m$ is homeomorphic to S^n , where $S^n \subseteq \mathbb{R}^{n+1}$ is the unit sphere, then (1.1) holds;
- (ii) if $S \subseteq \mathbb{R}^m$ is such that there exists a homeomorphism $f : B^n \rightarrow S$, where $B^n \subseteq \mathbb{R}^n$ is the unit ball, such that $f(S^{n-1})$ is a co-c.e. set, then (1.1) holds.

In the case $n = 1$, i.e. in the case when S is a topological circle or when S is a co-c.e. arc with computable endpoints, the preceding result has been generalized in [5] to computable metric spaces with the effective covering property and compact closed balls. Furthermore, by [6], the assumption of the effective covering property and compact closed balls can be replaced here by the weaker assumption that a computable metric space is locally computable.

In this paper we prove that this result holds for every $m \geq 1$, i.e. we prove that if (X, d, α) is a computable metric space which is locally computable, then

1998 ACM Subject Classification: F.1.1, F.4.1, G.0.

Key words and phrases: computable metric space, computable set, co-c.e. set, n -chain, spherical n -chain.

- (i) if S is a co-c.e. set in (X, d, α) homeomorphic to S^n , then S is computable in (X, d, α) ;
- (ii) if $f : B^n \rightarrow S$ is a homeomorphism, where S is co-c.e. in (X, d, α) , such that $f(S^{n-1})$ is also co-c.e., then S is computable.

In order to prove this, we use techniques similar to those in [5]. In Section 3 we examine the topological side of the problem. We define the notions of n -chain and spherical n -chain in a metric space. These notions play the same role as the notions of a chain and a circular chain play in the proof of the main results of [5]. However, the higher-dimensional aspect of the problem will require some deeper topological facts and we will rely here on a result proved in [4]. In Section 4 we include computability into consideration and we prove the main results of the paper: Theorem 4.8 and Theorem 4.9.

2. PRELIMINARIES

If X is a set, let $\mathcal{P}(X)$ denote the set of all subsets of X .

For $m \in \mathbb{N}$ let $\mathbb{N}_m = \{0, \dots, m\}$. For $n \geq 1$ let

$$\mathbb{N}_m^n = \{(x_1, \dots, x_n) \mid x_1, \dots, x_n \in \mathbb{N}_m\}.$$

We say that a function $\Phi : \mathbb{N}^k \rightarrow \mathcal{P}(\mathbb{N}^n)$ is **computable** if the function $\bar{\Phi} : \mathbb{N}^{k+n} \rightarrow \mathbb{N}$ defined by

$$\bar{\Phi}(x, y) = \chi_{\Phi(x)}(y),$$

$x \in \mathbb{N}^k, y \in \mathbb{N}^n$ is computable (i.e. recursive). Here $\chi_S : \mathbb{N}^n \rightarrow \{0, 1\}$ denotes the characteristic function of $S \subseteq \mathbb{N}^n$. A function $\Phi : \mathbb{N}^k \rightarrow \mathcal{P}(\mathbb{N}^n)$ is said to be **computably bounded** if there exists a computable function $\varphi : \mathbb{N}^k \rightarrow \mathbb{N}$ such that $\Phi(x) \subseteq \mathbb{N}_{\varphi(x)}^n$, for all $x \in \mathbb{N}^k$.

We say that a function $\Phi : \mathbb{N}^k \rightarrow \mathcal{P}(\mathbb{N}^n)$ is **c.c.b.** if Φ is computable and computably bounded.

Proposition 2.1.

- (1) If $\Phi, \Psi : \mathbb{N}^k \rightarrow \mathcal{P}(\mathbb{N}^n)$ are c.c.b. functions, then the sets $\{x \in \mathbb{N}^k \mid \Phi(x) = \Psi(x)\}$ and $\{x \in \mathbb{N}^k \mid \Phi(x) \subseteq \Psi(x)\}$ are decidable.
- (2) Let $\Phi : \mathbb{N}^k \rightarrow \mathcal{P}(\mathbb{N}^n)$ and $\Psi : \mathbb{N}^n \rightarrow \mathcal{P}(\mathbb{N}^m)$ be c.c.b. functions. Let $\Lambda : \mathbb{N}^k \rightarrow \mathcal{P}(\mathbb{N}^m)$ be defined by

$$\Lambda(x) = \bigcup_{z \in \Phi(x)} \Psi(z),$$

$x \in \mathbb{N}^k$. Then Λ is a c.c.b. function.

- (3) Let $\Phi : \mathbb{N}^k \rightarrow \mathcal{P}(\mathbb{N}^n)$ be c.c.b. and let $T \subseteq \mathbb{N}^n$ be c.e. Then the set $S = \{x \in \mathbb{N}^k \mid \Phi(x) \subseteq T\}$ is c.e. □

A function $F : \mathbb{N}^{k+1} \rightarrow \mathbb{Q}$ is called **computable** if there exist computable functions $a, b, c : \mathbb{N}^{k+1} \rightarrow \mathbb{N}$ such that

$$F(x) = (-1)^{c(x)} \frac{a(x)}{b(x) + 1}$$

for each $x \in \mathbb{N}^{k+1}$. A number $x \in \mathbb{R}$ is said to be **computable** if there exists a computable function $g : \mathbb{N} \rightarrow \mathbb{Q}$ such that $|x - g(i)| < 2^{-i}$ for each $i \in \mathbb{N}$.

By a **computable** function $\mathbb{N}^k \rightarrow \mathbb{R}$ we mean a function $f : \mathbb{N}^k \rightarrow \mathbb{R}$ for which there exists a computable function $F : \mathbb{N}^{k+1} \rightarrow \mathbb{Q}$ such that

$$|f(x) - F(x, i)| < 2^{-i}$$

for all $x \in \mathbb{N}^k$ and $i \in \mathbb{N}$.

In the following proposition we state some elementary facts about computable functions $\mathbb{N}^k \rightarrow \mathbb{R}$.

Proposition 2.2.

- (1) If $f, g : \mathbb{N}^k \rightarrow \mathbb{R}$ are computable, then $f + g, f - g : \mathbb{N}^k \rightarrow \mathbb{R}$ are computable.
- (2) If $f : \mathbb{N}^k \rightarrow \mathbb{R}$ and $F : \mathbb{N}^{k+1} \rightarrow \mathbb{R}$ are functions such that F is computable and $|f(x) - F(x, i)| < 2^{-i}$, $\forall x \in \mathbb{N}^k, \forall i \in \mathbb{N}$, then f is computable.
- (3) If $f : \mathbb{N}^{n+1} \rightarrow \mathbb{R}$ and $\varphi : \mathbb{N} \rightarrow \mathbb{N}$ are computable functions, then the function $g : \mathbb{N} \rightarrow \mathbb{R}$ defined by

$$g(l) = \max_{0 \leq j_1, \dots, j_n \leq \varphi(l)} f(l, j_1, \dots, j_n)$$

is computable.

- (4) If $f, g : \mathbb{N}^k \rightarrow \mathbb{R}$ are computable functions, then the set $\{x \in \mathbb{N}^k \mid f(x) > g(x)\}$ is c.e. □

A tuple (X, d, α) is said to be a **computable metric space** if (X, d) is a metric space and $\alpha : \mathbb{N} \rightarrow X$ is a sequence dense in (X, d) (i.e. a sequence whose range is dense in (X, d)) such that the function $\mathbb{N}^2 \rightarrow \mathbb{R}$,

$$(i, j) \mapsto d(\alpha_i, \alpha_j)$$

is computable (we use notation $\alpha = (\alpha_i)$).

If (X, d, α) is a computable metric space, then a sequence (x_i) in X is said to be **computable** in (X, d, α) if there exists a computable function $F : \mathbb{N}^2 \rightarrow \mathbb{N}$ such that

$$d(x_i, \alpha_{F(i,k)}) < 2^{-k}$$

for all $i, k \in \mathbb{N}$. A point $a \in X$ is said to be **computable** in (X, d, α) if the constant sequence a, a, \dots is computable.

Let (X, d, α) be a computable metric space. Let $q : \mathbb{N} \rightarrow \mathbb{Q}$ be some fixed computable function whose image is $\mathbb{Q} \cap \langle 0, \infty \rangle$ and let $\tau, \tau' : \mathbb{N} \rightarrow \mathbb{N}$ be some fixed computable functions such that $\{(\tau(i), \tau'(i)) \mid i \in \mathbb{N}\} = \mathbb{N}^2$. For $i \in \mathbb{N}$ we define

$$I_i = B(\alpha_{\tau(i)}, q_{\tau'(i)}), \quad \widehat{I}_i = \widehat{B}(\alpha_{\tau(i)}, q_{\tau'(i)}).$$

Here, for $x \in X$ and $r > 0$, we denote by $B(x, r)$ the open ball of radius r centered at x and by $\widehat{B}(x, r)$ the corresponding closed ball, i.e. $B(x, r) = \{y \in X \mid d(x, y) < r\}$, $\widehat{B}(x, r) = \{y \in X \mid d(x, y) \leq r\}$. For $A \subseteq X$ we will denote the closure of A by \overline{A} .

As a consequence of Proposition 2.2 we get the following corollary.

Corollary 2.3. *Let (X, d, α) be a computable metric space. The set $\{(k, i) \in \mathbb{N}^2 \mid \alpha_k \in I_i\}$ is c.e. □*

A closed subset S of (X, d) is said to be **computably enumerable** in (X, d, α) if

$$\{i \in \mathbb{N} \mid S \cap I_i \neq \emptyset\}$$

is a c.e. subset of \mathbb{N} . A closed subset S is said to be **co-computably enumerable** in (X, d, α) if there exists a computable function $f : \mathbb{N} \rightarrow \mathbb{N}$ such that

$$X \setminus S = \bigcup_{i \in \mathbb{N}} I_{f(i)}.$$

It is easy to see that these definitions do not depend on functions τ, τ' and q . We say that S is a **computable** set in (X, d, α) if S is both computably enumerable and co-computably enumerable ([2, 10]).

Let $\sigma : \mathbb{N}^2 \rightarrow \mathbb{N}$ and $\eta : \mathbb{N} \rightarrow \mathbb{N}$ be some fixed computable functions with the following property: $\{(\sigma(j, 0), \dots, \sigma(j, \eta(j))) \mid j \in \mathbb{N}\}$ is the set of all finite sequences in \mathbb{N} excluding the empty sequence, i.e. the set $\{(a_0, \dots, a_n) \mid n \in \mathbb{N}, a_0, \dots, a_n \in \mathbb{N}\}$. Such functions, for instance, can be defined using the Cantor pairing function. We use the following notation: $(j)_i$ instead of $\sigma(j, i)$ and \bar{j} instead of $\eta(j)$. Hence

$$\{((j)_0, \dots, (j)_{\bar{j}}) \mid j \in \mathbb{N}\}$$

is the set of all finite sequences in \mathbb{N} . For $j \in \mathbb{N}$ let $[j]$ be defined by

$$[j] = \{(j)_i \mid 0 \leq i \leq \bar{j}\}. \quad (2.1)$$

Note that the function $\mathbb{N} \rightarrow \mathcal{P}(\mathbb{N}), j \mapsto [j]$, is c.c.b.

Let (X, d, α) be a computable metric space. For $j \in \mathbb{N}$ we define

$$J_j = \bigcup_{i \in [j]} I_i, \quad \widehat{J}_j = \bigcup_{i \in [j]} \widehat{I}_i.$$

The sets J_j represent finite unions of rational balls and the sets \widehat{J}_j finite unions of closed rational balls.

Corollary 2.4. *Let (X, d, α) be a computable metric space. The set $\{(k, j) \in \mathbb{N}^2 \mid \alpha_k \in J_j\}$ is c.e.*

Proof. We have $\alpha_k \in J_j$ if and only if there exists $i \in \mathbb{N}$ such that $i \leq \bar{j}$ and $\alpha_k \in I_{(j)_i}$ and the claim follows from Corollary 2.3. \square

A computable metric space (X, d, α) has the **effective covering property** if the set

$$\{(w, j) \in \mathbb{N}^2 \mid \widehat{I}_w \subseteq J_j\}$$

is computably enumerable ([2]). It is not hard to see that this definition does not depend on the choice of the functions $q, \tau, \tau', \sigma, \eta$ which are necessary in the definitions of sets I_w and J_j .

For example, if $\alpha : \mathbb{N} \rightarrow \mathbb{R}^n$ is a computable function (in the sense that the component functions of α are computable) whose image is dense in \mathbb{R}^n and d is the Euclidean metric on \mathbb{R} , then $(\mathbb{R}^n, d, \alpha)$ is a computable metric space. A sequence (x_i) is computable in this computable metric space if and only if (x_i) is a computable sequence in \mathbb{R}^n and $(x_1, \dots, x_n) \in \mathbb{R}^n$ is a computable point in this space if and only if x_1, \dots, x_n are computable numbers. This computable metric space has the effective covering property (see e.g. [5]).

If (X, d, α) is a computable metric space, then a compact set K in (X, d) is said to be **computably compact** in (X, d, α) if K is computably enumerable in (X, d, α) and if the set $\{j \in \mathbb{N} \mid K \subseteq J_j\}$ is c.e. ([1]). A computable metric space (X, d, α) is **locally computable** ([1]) if for each compact set A in (X, d) there exists a computably compact set K in (X, d, α) such that $A \subseteq K$.

Let (X, d, α) be a computable metric space. A computable metric space (Y, d', β) is said to be a **subspace** of (X, d, α) if $Y \subseteq X$, $d' : Y \times Y \rightarrow \mathbb{R}$ is the restriction of $d : X \times X \rightarrow \mathbb{R}$ and β is a computable sequence in (X, d, α) .

The proofs of the following propositions can be found in [6].

Proposition 2.5. *Let (Y, d', β) be a subspace of a computable metric space (X, d, α) and let $S \subseteq Y$.*

- (i) *If S is co-c.e. in (X, d, α) , then S is co-c.e. in (Y, d', β) .*
- (ii) *If S is c.e. in (X, d, α) , then S is c.e. in (Y, d', β) . Conversely, if S is closed in (X, d) and c.e. in (Y, d', β) , then S is c.e. in (X, d, α) . \square*

Proposition 2.6. *Let (X, d, α) be a computable metric space and let K be a nonempty compact set in (X, d) . Then K is computably compact in (X, d, α) if and only if there exist a metric d' on K and a sequence β in K such that (K, d', β) is a subspace of (X, d, α) and (K, d', β) has the effective covering property. \square*

3. n -CHAINS AND SPHERICAL n -CHAINS

For $n \geq 1$ let

$$B^n = \{x \in \mathbb{R}^n \mid \|x\| \leq 1\}$$

and

$$S^{n-1} = \{x \in \mathbb{R}^n \mid \|x\| = 1\}.$$

A topological space X is called an n -**cell** if it is homeomorphic to B^n . We say that X is an n -**sphere** if it is homeomorphic to S^n .

By the **boundary sphere** of an n -cell E we mean the set $f(S^{n-1})$, where $f : B^n \rightarrow E$ is a homeomorphism. (Note that the boundary sphere of E , when E is a subspace of some topological space X , need not be equal to the topological boundary of E in X .) The definition of the boundary sphere does not depend on a particular homeomorphism $f : B^n \rightarrow E$. Namely, this is a consequence of the fact that each homeomorphism $B^n \rightarrow B^n$ maps S^{n-1} onto S^{n-1} (or equivalently $B^n \setminus S^{n-1}$ onto $B^n \setminus S^{n-1}$) which follows from the Invariance of domain theorem (see [8]): if $h : U \rightarrow \mathbb{R}^n$ is continuous and injective, where U is an open subset of \mathbb{R}^n , then $h(U)$ is open.

The result that we want to prove can now be restated in this way: if (X, d, α) is a computable metric space which is locally computable, then

- (1) each co-c.e. n -sphere is computable;
- (2) each co-c.e. n -cell whose boundary sphere is co-c.e. is computable.

Let us first note that it is enough to prove this result in the case when (X, d, α) is a computable metric space which has the effective covering property and compact closed balls. Namely, suppose that the result holds for such computable metric spaces and let (X, d, α) be a computable metric space which is locally computable. Let $S \subseteq X$ be a co-c.e. n -sphere. Then $S \subseteq K$, where K is computably compact in (X, d, α) . By Proposition 2.6 there exist d' and β such that (K, d', β) is a subspace of (X, d, α) and such that (K, d', β) has the effective covering property. By Proposition 2.5(i) S is co-c.e. in (K, d', β) and therefore S is computable in (K, d', β) . Proposition 2.5(ii) implies now that S is c.e. in (X, d, α) , hence S is computable in (X, d, α) . In the same way we get that each co-c.e. n -cell in (X, d, α) whose boundary sphere is co-c.e. is computable.

Let us observe how the statement (2) was proved in [5] in the case $n = 1$. Let E be a co-c.e. arc with computable endpoints a and b . For each $\varepsilon > 0$ there exists a finite sequence of open sets C_0, \dots, C_m such that

- (i) $E \subseteq C_0 \cup \dots \cup C_m$;
- (ii) $a \in C_0, b \in C_m$;

- (iii) $C_i \cap C_j = \emptyset$ for all i, j such that $|i - j| > 1$;
- (iv) each C_i is the finite union of rational balls, i.e. it is equal to some J_j ;
- (v) $\text{diam } C_i < \varepsilon$,

where $\text{diam } C_i$ denotes the diameter of the set C_i (see Figure 1). Since S is co-c.e. and (X, d, α) has the effective covering property and compact closed balls, it is possible to find effectively for each $k \in \mathbb{N}$ sets C_0, \dots, C_m with properties (i)–(v), where $\varepsilon = 2^{-k}$.




Figure 1.




Figure 2.

However, this means that we can effectively approximate S , namely properties (i)–(v) imply that $C_0 \cup \dots \cup C_m$ is a 2^{-k} -approximation of S in the following sense: for each $x \in E$ there exists $y \in C_0 \cup \dots \cup C_m$ such that $d(x, y) < 2^{-k}$ and for each $y \in C_0 \cup \dots \cup C_m$ there exists $x \in E$ such that $d(y, x) < 2^{-k}$. Using this fact we can prove that E is computable.

Why properties (i)–(v) imply that $C_0 \cup \dots \cup C_m$ is an 2^{-k} -approximation of S ? The fact that for each $x \in E$ there exists $y \in C_0 \cup \dots \cup C_m$ such that $d(x, y) < 2^{-k}$ follows trivially from (i). On the other hand, the fact that for each $y \in C_0 \cup \dots \cup C_m$ there exists $x \in E$ such that $d(y, x) < 2^{-k}$ can be easily deduced from (v) and the fact that

$$C_i \cap S \neq \emptyset \text{ for each } i \in \{0, \dots, m\}. \quad (3.1)$$

But why (3.1) holds? If we assume $C_i \cap E = \emptyset$ for some $i \in \{0, \dots, m\}$, then $0 < i < m$ and $C_0 \cup \dots \cup C_{i-1}$ and $C_{i+1} \cup \dots \cup C_m$ are two disjoint open sets (Figure 2.) which cover E and each of them intersects E which contradicts the fact that E is connected.

Suppose now that E is a 2-cell which is co-c.e. and whose boundary sphere is co-c.e. In order to prove that E is computable, we would like to proceed similarly as in the case of an arc. Naturally, in this case we are trying to find sets $C_{i,j}$, $0 \leq i, j \leq m$, which satisfy properties similar to properties (i)–(v) with basic difference that instead of (iii) we require

$$C_{i,j} \cap C_{i',j'} = \emptyset \text{ if } |i - i'| > 1 \text{ or } |j - j'| > 1. \quad (3.2)$$




Figure 3. The sets $C_{i,j}$ cover the 2-cell whose boundary sphere is the red curve

The main question here is what other properties we should require so that those properties imply

$$C_{i,j} \cap E \neq \emptyset \text{ for all } i, j \in \{0, \dots, m\}; \quad (3.3)$$

the fact (3.3) is important since we want to conclude that $\bigcup_{i,j} C_{i,j}$ approximates E in the same way as in the case of an arc.

If we suppose that i_0 and j_0 are such that $0 < i_0 < m$, $0 < j_0 < m$ and such that $C_{i_0, j_0} \cap E = \emptyset$, then we cannot conclude in general that E is covered by two disjoint open sets as in the case of an arc, but we can define the sets

$$U = \bigcup_{i < i_0} \bigcup_j C_{i,j}, \quad U' = \bigcup_{i > i_0} \bigcup_j C_{i,j},$$

$$V = \bigcup_{j < j_0} \bigcup_i C_{i,j}, \quad V' = \bigcup_{j > j_0} \bigcup_i C_{i,j},$$

and then these sets cover E and we have $U \cap U' = \emptyset$, $V \cap V' = \emptyset$. (See Figure 4. The missing set is C_{i_0, j_0} . The vertical blue sets are U and U' , the horizontal blue sets are V and V' .)




Figure 4.

Since E is homeomorphic to $I^2 = [0, 1] \times [0, 1]$, this raises the following question: is it possible to cover I^2 by open sets U , U' , V and V' so that $U \cap U' = \emptyset$, $V \cap V' = \emptyset$ and so that (see Figure 5.)

$$\{0\} \times [0, 1] \subseteq U, \quad \{1\} \times [0, 1] \subseteq U', \quad [0, 1] \times \{0\} \subseteq V, \quad [0, 1] \times \{1\} \subseteq V'?$$




Figure 5.

Let X be a topological space and let A, B, L be subsets of X . We say that L is a **partition between** A and B (see [4]) if there exist open sets U and W in X such that

$$A \subseteq U, B \subseteq W, U \cap W = \emptyset \text{ and } X \setminus L = U \cup W.$$

For $n \geq 1$ let

$$I^n = \{(x_1, \dots, x_n) \in \mathbb{R}^n \mid x_1, \dots, x_n \in [0, 1]\}.$$

For $n \geq 1$ and $i \in \{1, \dots, n\}$ let

$$A_i^{n,0} = \{(x_1, \dots, x_n) \in I^n \mid x_i = 0\},$$

$$A_i^{n,1} = \{(x_1, \dots, x_n) \in I^n \mid x_i = 1\}.$$

When the context is clear, we write A_i^0 and A_i^1 instead of $A_i^{n,0}$ and $A_i^{n,1}$. Let ∂I^n denote the boundary of I^n in \mathbb{R}^n , hence

$$\partial I^n = A_1^0 \cup \dots \cup A_n^0 \cup A_1^1 \cup \dots \cup A_n^1.$$

It is a well known fact that there is a homeomorphism $h : B^n \rightarrow I^n$ such that $h(S^{n-1}) = \partial I^n$. Hence if E is an n -cell, then there is a homeomorphism $f : I^n \rightarrow E$. In this case $f(\partial I^n)$ is the boundary sphere of E .

The following theorem can be found in [4] (Theorem 1.8.1).

Theorem 3.1. *Let $n \geq 1$. If L_i is a partition between A_i^0 and A_i^1 in I^n for $i \in \{1, \dots, n\}$, then $\bigcap_{i=1}^n L_i \neq \emptyset$. \square*

Corollary 3.2. *Let $n \geq 1$. Suppose U_1, \dots, U_n and V_1, \dots, V_n are open subsets of I^n such that*

$$U_i \cap A_i^1 = \emptyset, V_i \cap A_i^0 = \emptyset \text{ and } U_i \cap V_i = \emptyset$$

for all $i \in \{1, \dots, n\}$. Then $I^n \neq U_1 \cup \dots \cup U_n \cup V_1 \cup \dots \cup V_n$.

Proof. Suppose the opposite. Then $\{U_1, \dots, U_n, V_1, \dots, V_n\}$ is an open cover of I^n and let λ be its Lebesgue number. We can certainly find finitely many closed subsets B_1, \dots, B_l of I^n whose union is I^n and each of which has the diameter less than λ . Then each of the sets B_1, \dots, B_l is contained in some of the sets $U_1, \dots, U_n, V_1, \dots, V_n$.

For $i \in \{1, \dots, n\}$ we define F_i^0 to be the union of A_i^0 and all sets B_1, \dots, B_l which are subsets of U_i and F_i^1 to be the union of A_i^1 and all B_1, \dots, B_l which are subsets of V_i . Then $F_1^0, \dots, F_n^0, F_1^1, \dots, F_n^1$ are closed subsets of I^n , their union is I^n and for each $i \in \{1, \dots, n\}$ we have

$$A_i^0 \subseteq F_i^0, A_i^1 \subseteq F_i^1 \text{ and } F_i^0 \cap F_i^1 = \emptyset.$$

Let $i \in \{1, \dots, n\}$. Since F_i^0 and F_i^1 are closed and disjoint, there exist open sets W_i^0 and W_i^1 in I^n which are disjoint and such that $F_i^0 \subseteq W_i^0$, $F_i^1 \subseteq W_i^1$. Let $L_i = I^n \setminus (W_i^0 \cup W_i^1)$. Then L_i is a partition between A_i^0 and A_i^1 . We have

$$\bigcap_{i=1}^n L_i = I^n \setminus \bigcup_{i=1}^n (W_i^0 \cup W_i^1) \subseteq I^n \setminus \bigcup_{i=1}^n (F_i^0 \cup F_i^1) = \emptyset,$$

which is impossible by Theorem 3.1. \square

Corollary 3.3. *Let $n \geq 2$. Suppose U_1, \dots, U_{n-1} and V_1, \dots, V_{n-1} are open subsets of ∂I^n such that*

$$U_i \cap (A_i^{n,1} \cap A_n^{n,0}) = \emptyset, \quad V_i \cap (A_i^{n,0} \cap A_n^{n,0}) = \emptyset \quad \text{and} \quad U_i \cap V_i = \emptyset$$

for all $i \in \{1, \dots, n-1\}$. Let E be the union of all A_i^ρ such that $1 \leq i \leq n$, $\rho \in \{0, 1\}$, $(i, \rho) \neq (n, 0)$, i.e.

$$E = A_1^{n,0} \cup \dots \cup A_{n-1}^{n,0} \cup A_1^{n,1} \cup \dots \cup A_n^{n,1}.$$

Then E is not contained in the union $U_1 \cup \dots \cup U_{n-1} \cup V_1 \cup \dots \cup V_{n-1}$.




Figure 6.

Proof. See Figure 6. (case $n = 3$): the left and right blue sets are U_1 and U_2 respectively, and the left and right green sets are V_1 and V_2 respectively. In this figure E equals the union of vertical faces of the cube and the upper face of the cube.

Let $f : E \rightarrow I^{n-1}$ be defined by

$$f(x_1, \dots, x_n) = \left(\frac{1}{2} + \frac{1}{2x_n + 1} \left(x_1 - \frac{1}{2} \right), \dots, \frac{1}{2} + \frac{1}{2x_n + 1} \left(x_{n-1} - \frac{1}{2} \right) \right).$$

It is straightforward to check that f is bijective. Since E is compact and f clearly continuous, f is a homeomorphism. For each $i \in \{1, \dots, n-1\}$ we have

$$f(A_i^{n,0} \cap A_n^{n,0}) = A_i^{n-1,0}, \quad f(A_i^{n,1} \cap A_n^{n,0}) = A_i^{n-1,1}.$$

Suppose that $E \subseteq U_1 \cup \dots \cup U_{n-1} \cup V_1 \cup \dots \cup V_{n-1}$. Then

$$I^{n-1} = f(E \cap U_1) \cup \dots \cup f(E \cap U_{n-1}) \cup f(E \cap V_1) \cup \dots \cup f(E \cap V_n).$$

For each $i \in \{1, \dots, n-1\}$ the sets $f(E \cap U_i)$ and $f(E \cap V_i)$ are open in I^{n-1} , disjoint and

$$f(E \cap U_i) \cap A_i^{n-1,1} = \emptyset, \quad f(E \cap V_i) \cap A_i^{n-1,0} = \emptyset.$$

This is impossible by Corollary 3.2. \square

For $i \in \{1, \dots, n\}$ let

$$\partial_i^0 \mathbb{N}_m^n = \{(x_1, \dots, x_n) \in \mathbb{N}_m^n \mid x_i = 0\}, \quad \partial_i^1 \mathbb{N}_m^n = \{(x_1, \dots, x_n) \in \mathbb{N}_m^n \mid x_i = m\}$$

and let

$$\partial \mathbb{N}_m^n = \left(\bigcup_{1 \leq i \leq n} \partial_i^0 \mathbb{N}_m^n \right) \cup \left(\bigcup_{1 \leq i \leq n} \partial_i^1 \mathbb{N}_m^n \right).$$

Let X be a set, $n \geq 1$ and $m \in \mathbb{N}$. A function

$$C : \mathbb{N}_m^n \rightarrow \mathcal{P}(X)$$

is called an n -**chain** in X (of length m) if

$$C_{i_1, \dots, i_n} \cap C_{j_1, \dots, j_n} = \emptyset \quad (3.4)$$

for all $(i_1, \dots, i_n), (j_1, \dots, j_n) \in \mathbb{N}_m^n$ such that $|i_l - j_l| > 1$ for some $l \in \{1, \dots, n\}$. Here we use C_{i_1, \dots, i_n} to denote $C(i_1, \dots, i_n)$.

A **spherical** $(n-1)$ -**chain** in X (of length m) is a function

$$C : \partial \mathbb{N}_m^n \rightarrow \mathcal{P}(X)$$

such that (3.4) holds for all $(i_1, \dots, i_n), (j_1, \dots, j_n) \in \partial \mathbb{N}_m^n$ such that $|i_l - j_l| > 1$ for some $l \in \{1, \dots, n\}$.




Figure 7. A 2-chain and a spherical 1-chain

If $C : \mathbb{N}_m^n \rightarrow \mathcal{P}(X)$ is a function, we define its **boundary** ∂C as the restriction of C to $\partial \mathbb{N}_m^n$. For $i \in \{1, \dots, n\}$ and $\rho \in \{0, 1\}$ we define $\partial_i^\rho C$ as the restriction of C to $\partial_i^\rho \mathbb{N}_m^n$. Note: if C is an n -chain, then ∂C is a spherical $(n-1)$ -chain.

If $C : \partial \mathbb{N}_m^n \rightarrow \mathcal{P}(X)$ is a function and $i \in \{1, \dots, n\}$, $\rho \in \{0, 1\}$, we also use $\partial_i^\rho C$ to denote the restriction of the function C to $\partial_i^\rho \mathbb{N}_m^n$.

If (X, d) is a metric space, then we say that an n -chain $C = (C_{i_1, \dots, i_n})_{0 \leq i_1, \dots, i_n \leq m}$ in X is **open** if C_{i_1, \dots, i_n} is an open set in (X, d) for all $i_1, \dots, i_n \in \mathbb{N}_m$. We similarly define the notion of a **compact** n -chain in (X, d) and the notions of open spherical n -chain and compact spherical n -chain.

In general, if A is a set and $f : A \rightarrow \mathcal{P}(X)$ a function, we will denote by $\bigcup f$ the union $\bigcup_{a \in A} f(a)$ and we will say that f **covers** S , where $S \subseteq X$, if $S \subseteq \bigcup f$. If (X, d) is a metric space and $f(a)$ a nonempty bounded set for each $a \in A$, then we define $\text{mesh}(f)$ as the number

$$\text{mesh}(f) = \max_{a \in A} (\text{diam } f(a)).$$

Let $\varepsilon > 0$. A (spherical) n -chain C in a metric space (X, d) is said to be a (spherical) ε - n -**chain** if $\text{mesh}(C) < \varepsilon$.

A function $C : A \rightarrow \mathcal{P}(X)$, where $A \subseteq \mathbb{N}_m^n$, is said to be ε -**proper** if for all $(i_1, \dots, i_n), (j_1, \dots, j_n) \in A$ such that $|i_1 - j_1| \leq 1, \dots, |i_n - j_n| \leq 1$ there exist $x \in C_{i_1, \dots, i_n}$ and $y \in C_{j_1, \dots, j_n}$ such that $d(x, y) < \varepsilon$.

The proof of the following lemma is straightforward.

Lemma 3.4. *Let (X, d) be a metric space, $\varepsilon > 0$ and $A \subseteq \mathbb{N}_m^n$. Let $C, D : A \rightarrow \mathcal{P}(X)$ be such that $D(a) \neq \emptyset$ and $D(a) \subseteq C(a)$ for each $a \in A$. Suppose C is ε -proper and $\text{mesh}(C) < \varepsilon$. Then D is 3ε -proper. \square*

Lemma 3.5. *Let (X, d) be a metric space and let K be a compact (spherical) n -chain of length m in (X, d) . Suppose U_1, \dots, U_k are open sets. Then there exists an open (spherical) n -chain C of length m in (X, d) such that $K_a \subseteq C_a$ for all $a \in \mathbb{N}_m^n$ and such that $C_a \subseteq U_i$ whenever $i \in \{1, \dots, k\}$ is such that $K_a \subseteq U_i$. Moreover, if $\text{mesh}(K) < r$, we can choose C so that $\text{mesh}(C) < 2r$.*

Proof. If $S \subseteq X$ and $\varepsilon > 0$ let

$$S_\varepsilon = \bigcup_{s \in S} B(s, \varepsilon).$$

This is clearly an open set. If S is a compact set contained in some open set V , then there exists $\varepsilon > 0$ such that $S_\varepsilon \subseteq V$. Furthermore, if S and T are disjoint compact sets, then there exists $\varepsilon > 0$ such that $S_\varepsilon \cap T_\varepsilon = \emptyset$. It follows readily from this that there exists $\varepsilon > 0$ such that $C : \mathbb{N}_m^n \rightarrow \mathcal{P}(X)$ (or $C : \partial\mathbb{N}_m^n \rightarrow \mathcal{P}(X)$) defined by $C_a = (K_a)_\varepsilon$ is a desired n -chain (spherical n -chain). \square

Proposition 3.6. *Let $n \geq 2$. Suppose $f : \partial I^n \rightarrow S$ is a homeomorphism, where S is a subspace of a metric space (X, d) . Let $U_i^\rho, 1 \leq i \leq n, \rho \in \{0, 1\}$, be open sets in (X, d) such that*

$$f(A_i^\rho) \subseteq U_i^\rho$$

for all $i \in \{1, \dots, n\}$ and $\rho \in \{0, 1\}$. Then for each $\varepsilon > 0$ there exists an open spherical $\varepsilon - (n - 1)$ -chain C in (X, d) which is ε -proper, which covers S and such that

$$f(A_i^\rho) \subseteq \bigcup (\partial_i^\rho C) \subseteq U_i^\rho$$

for all $i \in \{1, \dots, n\}$ and $\rho \in \{0, 1\}$.

Proof. For $m \in \mathbb{N}$ let $D^m : \mathbb{N}_m^n \rightarrow \mathcal{P}(I^n)$ be defined by

$$D_{i_1, \dots, i_n}^m = \left[\frac{i_1}{m+1}, \frac{i_1+1}{m+1} \right] \times \dots \times \left[\frac{i_n}{m+1}, \frac{i_n+1}{m+1} \right].$$

Then D^m is a compact n -chain in I^n which covers I^n . Clearly for each $\varepsilon > 0$ there exists $m \in \mathbb{N}$ such that $\text{mesh}(D^m) < \varepsilon$. Note that for all $(i_1, \dots, i_n), (j_1, \dots, j_n) \in \mathbb{N}_m^n$ such that $|i_1 - j_1| \leq 1, \dots, |i_n - j_n| \leq 1$ we have

$$D_{i_1, \dots, i_n}^m \cap D_{j_1, \dots, j_n}^m \neq \emptyset.$$

We easily conclude from this that for each $\varepsilon > 0$ there exists $m \in \mathbb{N}$ such that $\text{mesh}(D^m) < \varepsilon$ and such that D^m is ε -proper.

The boundary ∂D^m is a spherical $(n - 1)$ -chain in I^n which covers ∂I^n .

For $m \in \mathbb{N}$ let $G^m : \partial\mathbb{N}_m^n \rightarrow \mathcal{P}(\partial I^n)$ be defined by

$$G^m(a) = (\partial D^m)(a) \cap \partial I^n.$$

Then G^m is a compact spherical $(n-1)$ -chain in ∂I^n which covers ∂I^n , moreover

$$A_i^\rho \subseteq \bigcup (\partial_i^\rho G^m)$$

for all $i \in \{1, \dots, n\}$ and $\rho \in \{0, 1\}$. Note that these sets need not be equal, however:

$$\text{for each } x \in \bigcup (\partial_i^\rho G^m) \text{ there exists } y \in A_i^\rho \text{ such that } d'(x, y) \leq \frac{1}{m+1}, \quad (3.5)$$

where d' is the Euclidean metric on \mathbb{R}^n . We also have that for each $\varepsilon > 0$ there exists $m \in \mathbb{N}$ such that $\text{mesh}(G^m) < \varepsilon$ and such that G^m is ε -proper.

For $m \in \mathbb{N}$ let $F^m : \partial \mathbb{N}_m^n \rightarrow \mathcal{P}(S)$ be defined by

$$F^m(a) = f(G^m(a)).$$

Then F^m is a compact spherical $(n-1)$ -chain in S which covers S and such that

$$f(A_i^\rho) \subseteq \bigcup (\partial_i^\rho F^m)$$

for all $i \in \{1, \dots, n\}$ and $\rho \in \{0, 1\}$.




Figure 8. D^3 , ∂D^3 , G^3 and F^3 (in the case $n = 2$)

The fact that f is uniformly continuous implies, together with (3.5), that for each $\varepsilon > 0$ there exists $m \in \mathbb{N}$ with the property that F^m is ε -proper, $\text{mesh}(F^m) < \varepsilon$ and with the property that for all $i \in \{1, \dots, n\}$, $\rho \in \{0, 1\}$ and $x \in \bigcup (\partial_i^\rho F^m)$ there exists $y \in f(A_i^\rho)$ such that $d(x, y) < \varepsilon$.

Let $\varepsilon > 0$. Using the fact that the sets A_i^ρ are compact and U_i^ρ are open, it is not hard to conclude now that there exists $m \in \mathbb{N}$ such that F^m is ε -proper, $\text{mesh}(F^m) < \frac{\varepsilon}{2}$ and

$$f(A_i^\rho) \subseteq \bigcup (\partial_i^\rho F^m) \subseteq U_i^\rho$$

for all $i \in \{1, \dots, n\}$ and $\rho \in \{0, 1\}$. Now we apply Lemma 3.5 to F and the sets U_i^ρ and we get an open spherical $\varepsilon - (n-1)$ -chain C in (X, d) which is ε -proper such that

$$f(A_i^\rho) \subseteq \bigcup (\partial_i^\rho C) \subseteq U_i^\rho$$

for all $i \in \{1, \dots, n\}$ and $\rho \in \{0, 1\}$. □

In the same way we prove the following proposition.

Proposition 3.7. *Let $n \geq 1$. Suppose $f : I^n \rightarrow E$ is a homeomorphism, where E is a subspace of a metric space (X, d) . Let U_i^ρ , $1 \leq i \leq n$, $\rho \in \{0, 1\}$, be open sets in (X, d) such that*

$$f(A_i^\rho) \subseteq U_i^\rho$$

for all $i \in \{1, \dots, n\}$ and $\rho \in \{0, 1\}$. Then for each $\varepsilon > 0$ there exists an open $\varepsilon - n$ -chain C in (X, d) which is ε -proper, which covers E and such that

$$f(A_i^\rho) \subseteq \bigcup (\partial_i^\rho C) \subseteq U_i^\rho$$

for all $i \in \{1, \dots, n\}$ and $\rho \in \{0, 1\}$.

If (X, d) is a metric space, then for nonempty subsets S and T of X we denote the number $\inf\{d(x, y) \mid x \in S, y \in T\}$ by $d(S, T)$.

The next proposition provides conditions under which a spherical $(n - 1)$ -chain approximates an $(n - 1)$ -sphere.

Proposition 3.8. *Let $f : \partial I^n \rightarrow S$ be a homeomorphism, where S is a subspace of a metric space (X, d) . Let W_i^ρ , $1 \leq i \leq n$, $\rho \in \{0, 1\}$, be open sets in (X, d) such that $W_i^0 \cap W_i^1 = \emptyset$ for all $i \in \{1, \dots, n\}$. Let $\varepsilon > 0$ be such that*

$$2\varepsilon < d(f(A_i^0), f(A_i^1)) \quad (3.6)$$

for each $i \in \{1, \dots, n\}$. Suppose C is an open spherical $\varepsilon - (n - 1)$ -chain in (X, d) of length m which is ε -proper, which covers S and suppose that

$$f(A_i^\rho) \subseteq W_i^\rho, \bigcup (\partial_i^\rho C) \subseteq W_i^\rho$$

for all $i \in \{1, \dots, n\}$ and $\rho \in \{0, 1\}$. Then for each $x \in \bigcup C$ there exists $y \in S$ such that $d(x, y) < 3\varepsilon$.

Proof. It is enough to prove the following: for each $(p_1, \dots, p_n) \in \partial \mathbb{N}_m^n$ with the property that $p_k \in \{0, m\}$ for exactly one $k \in \{1, \dots, n\}$ the set C_{p_1, \dots, p_n} intersects S . Namely, if this holds, then for each $(q_1, \dots, q_n) \in \partial \mathbb{N}_m^n$ there exists $(p_1, \dots, p_n) \in \partial \mathbb{N}_m^n$ such that $|q_1 - p_1| \leq 1, \dots, |q_n - p_n| \leq 1$ and such that $C_{p_1, \dots, p_n} \cap S \neq \emptyset$. Since C is an $\varepsilon - (n - 1)$ -chain and ε -proper, we now easily get that for each $x \in \bigcup C$ there exists $y \in S$ such that $d(x, y) < 3\varepsilon$.

Suppose the opposite, that there exists $(p_1, \dots, p_n) \in \partial \mathbb{N}_m^n$ such that $p_k \in \{0, m\}$ for exactly one $k \in \{1, \dots, n\}$ and such that $C_{p_1, \dots, p_n} \cap S = \emptyset$. We may assume $p_n = m$ (all other cases can be reduced to this one if we modify C and f by interchange of appropriate coordinates). It follows $0 < p_1 < m, \dots, 0 < p_{n-1} < m$.

For $i \in \{1, \dots, n - 1\}$ we define the set U_i as the union of all sets of the following form:

$$C_{j_1, \dots, j_{i-1}, l, j_{i+1}, \dots, j_{n-1}, m}, \text{ where } l < p_i; \quad (3.7)$$

$$C_{j_1, \dots, j_{i-1}, 0, j_{i+1}, \dots, j_n}; \quad (3.8)$$

$$C_{j_1, \dots, j_{n-1}, 0}, \text{ where this set is such that it intersects } f(A_i^0). \quad (3.9)$$

Furthermore, let V_i be the union of all sets of the following form:

$$C_{j_1, \dots, j_{i-1}, l, j_{i+1}, \dots, j_{n-1}, m}, \text{ where } l > p_i; \quad (3.10)$$

$$C_{j_1, \dots, j_{i-1}, m, j_{i+1}, \dots, j_n}; \quad (3.11)$$

$$C_{j_1, \dots, j_{n-1}, 0}, \text{ where this set is such that it intersects } f(A_i^1). \quad (3.12)$$

Let $i \in \{1, \dots, n - 1\}$. The sets U_i and V_i are open and it is straightforward to check that they are disjoint. (Figures 9. and 10. show C in case $n = 3$ and $m = 5$; the red set is C_{p_1, p_2, p_3} , in this case $C_{2, 2, 5}$, the blue sets in Figure 9. are U_1 and V_1 , the blue sets in Figure 10. are U_2 and V_2 . For example, note that in Figure 11. the black set is $C_{3, 0, 2}$, the red set is $C_{5, 0, 1}$ and the blue set is $C_{5, 0, 5}$.)




Figure 9.




Figure 10.




Figure 11.

We also have

$$U_i \cap f(A_i^1 \cap A_n^0) = \emptyset. \quad (3.13)$$

Otherwise, the set $f(A_i^1 \cap A_n^0) = f(A_i^1) \cap f(A_n^0)$ would intersect one of the sets in (3.7), (3.8) or (3.9). The sets in (3.7) are contained in W_n^1 which is disjoint with $f(A_n^0)$. The sets in (3.8) are contained in W_i^0 which is disjoint with $f(A_i^1)$. Finally, $f(A_i^1)$ cannot intersect a set in (3.9) since (3.6) holds. In the same way we get

$$V_i \cap f(A_i^0 \cap A_n^0) = \emptyset. \quad (3.14)$$

Let

$$\Omega = U_1 \cup \dots \cup U_{n-1} \cup V_1 \cup \dots \cup V_{n-1}.$$

Let $i \in \{1, \dots, n\}$ and $\rho \in \{0, 1\}$ such that $(i, \rho) \neq (n, 0)$. We claim that

$$f(A_i^\rho) \subseteq \Omega. \quad (3.15)$$

Suppose that there exists $x \in f(A_i^\rho)$ such that $x \notin \Omega$. Since C covers S , there exists $(j_1, \dots, j_n) \in \partial\mathbb{N}_m^n$ such that

$$x \in C_{j_1, \dots, j_n}.$$

We have $(j_1, \dots, j_n) \neq (p_1, \dots, p_n)$ since $C_{p_1, \dots, p_n} \cap S = \emptyset$. So, if $j_n = m$, then C_{j_1, \dots, j_n} must be one of the sets in (3.7) or (3.10). But this is impossible since $x \notin \Omega$. So $j_n < m$. Now, if $j_n > 0$, then C_{j_1, \dots, j_n} is one of the sets in (3.8) or (3.11), impossible. Therefore $j_n = 0$.

We have $C_{j_1, \dots, j_n} \cap f(A_i^\rho) \neq \emptyset$ and this also yields to contradiction. Namely, if $i < n$, then C_{j_1, \dots, j_n} is one of the sets in (3.9) or (3.12). And if $i = n$, then $\rho = 1$ and

$$f(A_n^1) \subseteq W_n^1, \quad C_{j_1, \dots, j_n} \subseteq \bigcup (\partial_n^0 C) \subseteq W_n^0,$$

which is impossible since $W_n^0 \cap W_n^1 = \emptyset$. Hence (3.15) holds.

Let

$$E = A_1^0 \cup \dots \cup A_{n-1}^0 \cup A_1^1 \cup \dots \cup A_n^1.$$

For each $i \in \{1, \dots, n-1\}$ the sets $f^{-1}(U_i)$ and $f^{-1}(V_i)$ are open in ∂I^n , they are disjoint, by (3.13) and (3.14)

$$f^{-1}(U_i) \cap (A_i^1 \cap A_n^0) = \emptyset, \quad f^{-1}(V_i) \cap (A_i^0 \cap A_n^0) = \emptyset$$

and by (3.15)

$$E \subseteq f^{-1}(U_1) \cup \dots \cup f^{-1}(U_{n-1}) \cup f^{-1}(V_1) \cup \dots \cup f^{-1}(V_{n-1}).$$

This is impossible by Corollary 3.3. \square

The next proposition provides conditions under which an n -chain approximates an n -cell.

Proposition 3.9. *Let $f : I^n \rightarrow E$ be a homeomorphism, where E is a subspace of a metric space (X, d) . Suppose C is an open ε - n -chain in (X, d) of length m which is ε -proper, which covers E and such that ∂C covers $f(\partial I^n)$ and suppose that W_i^ρ , $1 \leq i \leq n$, $\rho \in \{0, 1\}$, are open sets in (X, d) such that*

$$2\varepsilon < d(W_i^0, W_i^1),$$

$$f(A_i^\rho) \subseteq W_i^\rho \text{ and } \bigcup (\partial_i^\rho C) \subseteq W_i^\rho$$

for all $i \in \{1, \dots, n\}$ and $\rho \in \{0, 1\}$. Then for each $x \in \bigcup C$ there exists $y \in E$ such that $d(x, y) < 7\varepsilon$.

Proof. It is enough to prove that for each $(p_1, \dots, p_n) \in \mathbb{N}_m^n$ with the property that $1 < p_k < m-1$ for each $k \in \{1, \dots, n\}$ the set C_{p_1, \dots, p_n} intersects E . Why is it enough to prove this? Suppose that this fact holds. Assume that $m \geq 4$. Let $x \in \bigcup C$. Then $x \in C_{q_1, \dots, q_n}$ for some $(q_1, \dots, q_n) \in \mathbb{N}_m^n$. For $i \in \{1, \dots, n\}$ let

$$q'_i = \begin{cases} 1 & \text{if } q_i = 0, \\ m-1 & \text{if } q_i = m, \\ q_i & \text{otherwise.} \end{cases}$$

Then $|q_1 - q'_1| \leq 1, \dots, |q_n - q'_n| \leq 1$ and $1 \leq q'_i \leq m-1$ for each $i \in \{1, \dots, n\}$. Since C is ε -proper, there exist $x' \in C_{q_1, \dots, q_n}$ and $x'' \in C_{q'_1, \dots, q'_n}$ such that $d(x', x'') < \varepsilon$. Now, let the numbers p_1, \dots, p_n be defined by

$$p_i = \begin{cases} 2 & \text{if } q'_i = 1, \\ m-2 & \text{if } q'_i = m-1, \\ q'_i & \text{otherwise,} \end{cases}$$

$i \in \{1, \dots, n\}$. We have $|q'_1 - p_1| \leq 1, \dots, |q'_n - p_n| \leq 1$ and therefore there exist $y'' \in C_{q'_1, \dots, q'_n}$ and $y' \in C_{p_1, \dots, p_n}$ such that $d(y'', y') < \varepsilon$. Clearly $1 < p_i < m-1$ for each $i \in \{1, \dots, n\}$ and therefore there exists $y \in C_{p_1, \dots, p_n} \cap E$. Using the fact that the diameters of the sets C_{q_1, \dots, q_n} , $C_{q'_1, \dots, q'_n}$ and C_{p_1, \dots, p_n} are less than ε , we obtain

$$d(x, y) \leq d(x, x') + d(x', x'') + d(x'', y'') + d(y'', y') + d(y', y) < 5\varepsilon.$$

If $m \leq 3$, then for all $(a_1, \dots, a_n), (b_1, \dots, b_n) \in \mathbb{N}_m^n$ we have, for each $i \in \{1, \dots, n\}$, that $a_i, b_i \in \{0, \dots, m\} \subseteq \{0, 1, 2, 3\}$ and therefore there exist $c_i, d_i \in \{0, \dots, m\}$ such that $|a_i - c_i| \leq 1, |c_i - d_i| \leq 1, |d_i - b_i| \leq 1$ which, together with the fact that C is ε -proper, implies that there exist $x \in C_{a_1, \dots, a_n}$, $x', y' \in C_{c_1, \dots, c_n}$, $y'', z'' \in C_{d_1, \dots, d_n}$ and $z \in C_{b_1, \dots, b_n}$

such that $d(x, x') < \varepsilon$, $d(y', y'') < \varepsilon$ and $d(z'', z) < \varepsilon$. It follows $d(x, z) < 5\varepsilon$. This proves that $d(C_a, C_b) < 5\varepsilon$ for all $a, b \in \mathbb{N}_m^n$. Since E is nonempty and contained in $\bigcup C$, there exists $b \in \mathbb{N}_m^n$ such that $C_b \cap E \neq \emptyset$. It follows $d(C_a, E) < 6\varepsilon$ for each $a \in \mathbb{N}_m^n$ and therefore for each $x \in \bigcup C$ we have $d(x, E) < 7\varepsilon$.

So, let $(p_1, \dots, p_n) \in \mathbb{N}_m^n$ be such that $1 < p_k < m-1$ for each $k \in \{1, \dots, n\}$. We want to prove that $C_{p_1, \dots, p_n} \cap E \neq \emptyset$. Suppose $C_{p_1, \dots, p_n} \cap E = \emptyset$.

For $i \in \{1, \dots, n\}$ let U_i be the union of all sets C_{j_1, \dots, j_n} such that

$$j_i < p_i \text{ and } 1 < j_k < m-1 \text{ for all } k \in \{1, \dots, n\} \quad (3.16)$$

or

$$j_i \in \{0, 1\}. \quad (3.17)$$

Let V_i be the union of all sets C_{j_1, \dots, j_n} such that

$$j_i > p_i \text{ and } 1 < j_k < m-1 \text{ for all } k \in \{1, \dots, n\}$$

or

$$j_i \in \{m-1, m\}.$$

For each $i \in \{1, \dots, n\}$ the sets U_i and V_i are open and disjoint. Note that every C_{j_1, \dots, j_n} , where $(j_1, \dots, j_n) \neq (p_1, \dots, p_n)$, is contained in some U_i or V_i . Therefore

$$E \subseteq U_1 \cup \dots \cup U_n \cup V_1 \cup \dots \cup V_n. \quad (3.18)$$

Let $i \in \{1, \dots, n\}$. We prove now that

$$U_i \cap f(A_i^1) = \emptyset. \quad (3.19)$$

Suppose the opposite, that $U_i \cap f(A_i^1) \neq \emptyset$. It follows from the definition of U_i that there exist $j_1, \dots, j_n \in \mathbb{N}_m$ such that (3.16) or (3.17) hold and such that $C_{j_1, \dots, j_n} \cap f(A_i^1) \neq \emptyset$. However, if (3.16) holds, then $1 < j_k < m-1$ for all $k \in \{1, \dots, n\}$ which implies that C_{j_1, \dots, j_n} is disjoint with $\bigcup(\partial C)$. But we have the assumption that ∂C covers $f(\partial I^n)$ and this implies that C_{j_1, \dots, j_n} is disjoint with $f(\partial I^n)$ which is impossible. Therefore, (3.16) does not hold which means that (3.17) holds. Hence we have

$$C_{j_1, \dots, j_n} \cap f(A_i^1) \neq \emptyset \text{ and } j_i \in \{0, 1\}.$$

This, together with $\bigcup(\partial_i^0 C) \subseteq W_i^0$ and $f(A_i^1) \subseteq W_i^1$, implies $d(W_i^0, W_i^1) < 2\varepsilon$ (namely, if $j_i = 0$, then $W_i^0 \cap W_i^1 \neq \emptyset$, and if $j_i = 1$, then $d(C_{j_1, \dots, j_n}, W_i^0) < \varepsilon$ since C is ε -proper and this implies $d(W_i^0, W_i^1) < 2\varepsilon$). A contradiction. Hence (3.19) holds. In the same way we get

$$V_i \cap f(A_i^0) = \emptyset. \quad (3.20)$$

For each $i \in \{1, \dots, n\}$ the sets $f^{-1}(U_i)$ and $f^{-1}(V_i)$ are open in I^n and disjoint. By (3.18)

$$I^n = f^{-1}(U_1) \cup \dots \cup f^{-1}(U_n) \cup f^{-1}(V_1) \cup \dots \cup f^{-1}(V_n),$$

and by (3.19) and (3.20)

$$f^{-1}(U_i) \cap A_i^1 = \emptyset, \quad f^{-1}(V_i) \cap A_i^0 = \emptyset.$$

This is impossible by Corollary 3.2. □

4. COMPUTABILITY OF CO-C.E. SPHERES AND CELLS

Let $n \geq 1$. A **finite n -sequence** in \mathbb{N} is any function of the form

$$\{0, \dots, m\}^n \rightarrow \mathbb{N}.$$

Recall that any finite sequence i_0, \dots, i_m in \mathbb{N} is of the form $(j)_0, \dots, (j)_{\bar{j}}$ for some $j \in \mathbb{N}$. Let $f : \mathbb{N}^n \rightarrow \mathbb{N}$ be some computable injection and let τ and τ' be the functions from the section Preliminaries. We define $\Sigma : \mathbb{N}^{n+1} \rightarrow \mathbb{N}$ by

$$\Sigma(i, j_1, \dots, j_n) = (\tau(i))_{f(j_1, \dots, j_n)}.$$

Then for any finite n -sequence a in \mathbb{N} there exists $i \in \mathbb{N}$ such that a equals the function

$$\begin{aligned} \{0, \dots, \tau'(i)\}^n &\rightarrow \mathbb{N}, \\ (j_1, \dots, j_n) &\mapsto \Sigma(i, j_1, \dots, j_n). \end{aligned}$$

We will use the following notation: \widehat{i} instead of $\tau'(i)$ and, for $n \geq 2$, $(i)_{j_1, \dots, j_n}$ instead of $\Sigma(i, j_1, \dots, j_n)$.

Let (X, d, α) be a computable metric space. For $l \in \mathbb{N}$ let \mathcal{H}_l be the finite n -sequence of sets in X defined by

$$\mathcal{H}_l = \left(J_{(l)_{j_1, \dots, j_n}} \right)_{0 \leq j_1, \dots, j_n \leq \widehat{l}}$$

(i.e. \mathcal{H}_l is the function $\{0, \dots, \widehat{l}\}^n \rightarrow \mathcal{P}(X)$ which maps (j_1, \dots, j_n) to $J_{(l)_{j_1, \dots, j_n}}$).

For $l \in \mathbb{N}$ let $\widehat{\mathcal{H}}_l$ be defined by

$$\widehat{\mathcal{H}}_l = \left(\widehat{J}_{(l)_{j_1, \dots, j_n}} \right)_{0 \leq j_1, \dots, j_n \leq \widehat{l}}.$$

In Euclidean space \mathbb{R}^n we can effectively calculate the diameter of the finite union of rational balls. However, in a general computable metric space the function $\mathbb{N} \rightarrow \mathbb{R}$, $j \mapsto \text{diam}(J_j)$, need not be computable. For that reason we are going to use the notion of the formal diameter. Let (X, d) be a metric space and $x_0, \dots, x_k \in X$, $r_0, \dots, r_k \in \mathbb{R}_+$. The **formal diameter** associated to the finite sequence $(x_0, r_0), \dots, (x_k, r_k)$ is the number $D \in \mathbb{R}$ defined by

$$D = \max_{0 \leq v, w \leq k} d(x_v, x_w) + 2 \max_{0 \leq v \leq k} r_v.$$

Let (X, d, α) be a computable metric space. We define the function $\text{fdiam} : \mathbb{N} \rightarrow \mathbb{R}$ in the following way. For $j \in \mathbb{N}$ the number $\text{fdiam}(j)$ is the formal diameter associated to the finite sequence

$$\left(\alpha_{\tau((j)_0)}, q_{\tau'((j)_0)} \right), \dots, \left(\alpha_{\tau((j)_{\bar{j}})}, q_{\tau'((j)_{\bar{j}})} \right).$$

We have the following proposition (for the proof see [5]).

Proposition 4.1. *Let (X, d, α) be a computable metric space.*

- (1) *For all $j \in \mathbb{N}$, $\text{diam}(\widehat{J}_j) \leq \text{fdiam}(j)$.*
- (2) *$\text{fdiam} : \mathbb{N} \rightarrow \mathbb{R}$ is a computable function.*
- (3) *Let S be a compact subset of (X, d) , $r \in \mathbb{R}_+$ and C_0, \dots, C_m a finite sequence of open sets which covers S and such that $\text{diam}(C_i) < r$ for each $i \in \{0, \dots, m\}$. Then there exist $j_0, \dots, j_m \in \mathbb{N}$ such that the finite sequence of sets J_{j_0}, \dots, J_{j_m} covers S , $\widehat{J}_{j_i} \subseteq C_i$ and $\text{fdiam}(j_i) < r$ for each $i \in \{0, \dots, m\}$.*

Let the function $\text{fmesh} : \mathbb{N} \rightarrow \mathbb{R}$ be defined by

$$\text{fmesh}(l) = \max_{0 \leq j_1, \dots, j_n \leq \widehat{l}} \text{fdiam}((l)_{j_1, \dots, j_n}).$$

It is immediate from Proposition 4.1 and Proposition 2.2 that fmesh is a computable function.

Proposition 4.2. *Let (X, d, α) be a computable metric space. The sets*

$$\Omega = \{(l, k) \in \mathbb{N}^2 \mid \mathcal{H}_l \text{ is } 2^{-k} \text{ - proper}\}$$

and

$$\Omega' = \{(l, k) \in \mathbb{N}^2 \mid \partial \mathcal{H}_l \text{ is } 2^{-k} \text{ - proper}\}$$

are c.e.

Proof. Let $\Phi : \mathbb{N}^2 \rightarrow \mathcal{P}(\mathbb{N}^{2n+2})$ be defined in the following way. For $l, k \in \mathbb{N}$ let $\Phi(l, k)$ be the set of all

$$(l, k, i_1, \dots, i_n, j_1, \dots, j_n)$$

such that $i_1, \dots, i_n, j_1, \dots, j_n \in \mathbb{N}_{\widehat{l}}$ and $|i_1 - j_1| \leq 1, \dots, |i_n - j_n| \leq 1$. Then Φ is c.c.b. On the other hand, let S be the set of all $(l, k, i_1, \dots, i_n, j_1, \dots, j_n)$ for which there exists $x \in J_{(l)_{i_1, \dots, i_n}}$ and $y \in J_{(l)_{j_1, \dots, j_n}}$ such that $d(x, y) < 2^{-k}$. This is equivalent to the fact that there exist $p, q \in \mathbb{N}$ such that

$$\alpha_p \in J_{(l)_{i_1, \dots, i_n}}, \alpha_q \in J_{(l)_{j_1, \dots, j_n}} \text{ and } d(\alpha_p, \alpha_q) < 2^{-k} \quad (4.1)$$

The set T of all $(l, k, i_1, \dots, i_n, j_1, \dots, j_n, p, q)$ such that (4.1) holds is c.e. by Corollary 2.4 and Proposition 2.2. Therefore S is c.e. Since

$$\Omega = \{(l, k) \mid \Phi(l, k) \subseteq S\}$$

we have that Ω is c.e. by Proposition 2.1. We similarly get that Ω' is c.e. \square

Lemma 4.3. *Let (X, d, α) be a computable metric space. There exists a computable function $\zeta : \mathbb{N} \rightarrow \mathbb{N}$ such that $J_{\zeta(l)} = \bigcup \mathcal{H}_l$ for each $l \in \mathbb{N}$. There exists a computable function $\zeta' : \mathbb{N} \rightarrow \mathbb{N}$ such that $J_{\zeta'(l)} = \bigcup (\partial \mathcal{H}_l)$ for each $l \in \mathbb{N}$. Furthermore, for all $i \in \{1, \dots, m\}$ and $\rho \in \{0, 1\}$ there exists a computable function $\zeta'' : \mathbb{N} \rightarrow \mathbb{N}$ such that $J_{\zeta''(l)} = \bigcup (\partial_i^\rho \mathcal{H}_l)$ for each $l \in \mathbb{N}$. Similar statements hold for \widehat{J}_j and $\widehat{\mathcal{H}}_l, \partial \widehat{\mathcal{H}}_l, \partial_i^\rho \widehat{\mathcal{H}}_l$.*

Proof. It is enough to prove the following: if $\Phi : \mathbb{N} \rightarrow \mathcal{P}(\mathbb{N}^n)$ and $\Psi : \mathbb{N}^n \rightarrow \mathcal{P}(\mathbb{N})$ are c.c.b. functions such that $\Phi(l) \neq \emptyset$ and $\Psi(a) \neq \emptyset$ for all $l \in \mathbb{N}$ and $a \in \mathbb{N}^n$, then there exists a computable function $\zeta : \mathbb{N} \rightarrow \mathbb{N}$ such that

$$J_{\zeta(l)} = \bigcup_{a \in \Phi(l)} \bigcup_{i \in \Psi(a)} I_i. \quad (4.2)$$

However, if Φ and Ψ are such functions, by Proposition 2.1 there exists a c.c.b. function $\Lambda : \mathbb{N} \rightarrow \mathcal{P}(\mathbb{N})$ such that

$$\bigcup_{i \in \Lambda(l)} I_i = \bigcup_{a \in \Phi(l)} \bigcup_{i \in \Psi(a)} I_i.$$

For each $l \in \mathbb{N}$ there exists $j \in \mathbb{N}$ such that $\Lambda(l) = [j]$ (recall definition (2.1)). Since the set $S = \{(l, j) \mid \Lambda(l) = [j]\}$ is computable (Proposition 2.1) and for each $l \in \mathbb{N}$ there exists $j \in \mathbb{N}$ such that $(l, j) \in S$, there exists a computable function $\zeta : \mathbb{N} \rightarrow \mathbb{N}$ such that $(l, \zeta(l)) \in S$ for each $l \in \mathbb{N}$. It follows (4.2). \square

The proof of the following proposition can be found in [5].

Proposition 4.4. *Let (X, d, α) be a computable metric space which has the effective covering property and compact closed balls.*

- (1) *The set $\{(i, j) \in \mathbb{N}^2 \mid \widehat{J}_i \subseteq J_j\}$ is c.e.*
- (2) *Let S be a co-c.e. set in (X, d, α) which is compact. Then the set $\{j \in \mathbb{N} \mid S \subseteq J_j\}$ is c.e. \square*

Corollary 4.5. *Let (X, d, α) be a computable metric space which has the effective covering property and compact closed balls and let S be a co-c.e. set in (X, d, α) which is compact. Then the sets*

$$\{l \in \mathbb{N} \mid \mathcal{H}_l \text{ covers } S\} \text{ and } \{l \in \mathbb{N} \mid \partial \mathcal{H}_l \text{ covers } S\}$$

are c.e.

Proof. This follows from Lemma 4.3 and Proposition 4.4. \square

The following proposition can be proved in the same way as Proposition 32 in [5].

Proposition 4.6. *Let (X, d, α) be a computable metric space which has the effective covering property and compact closed balls. The sets $\Omega = \{l \in \mathbb{N} \mid \widehat{\mathcal{H}}_l \text{ is an } n\text{-chain}\}$ and $\Omega' = \{l \in \mathbb{N} \mid \partial \widehat{\mathcal{H}}_l \text{ is a spherical } (n-1)\text{-chain}\}$ are computably enumerable. \square*

The following lemma can be proved similarly as Lemma 14 in [5].

Lemma 4.7. *Let (X, d, α) be a computable metric space. Let S be a compact set in this space such that there exists a computable function $f : \mathbb{N} \rightarrow \mathbb{N}$ with the property that for each $k \in \mathbb{N}$ the following holds:*

$$S \subseteq J_{f(k)} \text{ and for each } x \in J_{f(k)} \text{ there exists } y \in S \text{ such that } d(x, y) < 2^{-k}.$$

Then S is computable. \square

Theorem 4.8. *Let (X, d, α) be a computable metric space which is locally computable. Let S be an $(n-1)$ -sphere in (X, d) and suppose S is co-c.e. in (X, d, α) . Then S is computable.*

Proof. As we have seen, we may assume that (X, d, α) has compact closed balls and the effective covering property. Let $f : \partial I^n \rightarrow S$ be a homeomorphism. Choose sets W_i^ρ , $1 \leq i \leq n$, $\rho \in \{0, 1\}$, so that each of these sets is a finite union of rational balls (i.e. of the form J_j) and so that

$$W_i^0 \cap W_i^1 = \emptyset \text{ and } f(A_i^\rho) \subseteq W_i^\rho$$

for all $i \in \{1, \dots, n\}$ and $\rho \in \{0, 1\}$.

Let $k_0 \in \mathbb{N}$ be such that

$$2 \cdot 2^{-k_0} < d(f(A_i^0), f(A_i^1))$$

for each $i \in \{1, \dots, n\}$.

By Proposition 3.6 for each $\varepsilon > 0$ there exists an open spherical $\varepsilon - (n-1)$ -chain C in (X, d) which is ε -proper, which covers S and such that

$$f(A_i^\rho) \subseteq \bigcup (\partial_i^\rho C) \subseteq W_i^\rho$$

for all $i \in \{1, \dots, n\}$ and $\rho \in \{0, 1\}$.

From this, Lemma 3.4 and Proposition 4.1 we conclude that for each $k \in \mathbb{N}$ there exists $l \in \mathbb{N}$ with the following properties:

$$\partial \widehat{\mathcal{H}}_l \text{ is a spherical } (n-1)\text{-chain,} \quad (4.3)$$

$$\partial \mathcal{H}_l \text{ covers } S, \quad (4.4)$$

$$\text{fmesh}(l) < 2^{-(k+k_0)}, \quad (4.5)$$

$$\partial \mathcal{H}_l \text{ is } 2^{-(k+k_0)} \text{ - proper} \quad (4.6)$$

and

$$\bigcup \left(\partial_i^\rho \widehat{\mathcal{H}}_l \right) \subseteq W_i^\rho \quad (4.7)$$

for all $i \in \{1, \dots, n\}$ and $\rho \in \{0, 1\}$.

Let Ω be the set of all (k, l) such that (4.3), (4.4), (4.5), (4.6) and (4.7) hold. Then Ω is c.e., which follows from Proposition 4.6, Corollary 4.5, Proposition 4.2, Lemma 4.3, Proposition 4.4(1) and the fact that fmesh is a computable function. The fact that Ω is c.e. and the fact that for each $k \in \mathbb{N}$ there exists $l \in \mathbb{N}$ such that $(k, l) \in \Omega$ imply that there exists a computable function $g : \mathbb{N} \rightarrow \mathbb{N}$ such that $(k, g(k)) \in \Omega$ for each $k \in \mathbb{N}$.

Let $k \in \mathbb{N}$. By Proposition 3.8 for each $x \in \bigcup (\partial \mathcal{H}_{g(k)})$ there exists $y \in S$ such that $d(x, y) < 3 \cdot 2^{-k}$. Now Lemma 4.3 and Lemma 4.7 imply that S is computable. \square

Theorem 4.9. *Let (X, d, α) be a computable metric space which is locally computable. Let E be an n -cell in (X, d) and suppose E and the boundary sphere of E are co-c.e. in (X, d, α) . Then E is computable.*

Proof. We proceed in a similar way as in the proof of Theorem 4.8. First, we may assume that (X, d, α) has compact closed balls and the effective covering property. Let $f : I^n \rightarrow E$ be a homeomorphism. Let $S = f(\partial I^n)$. Choose sets W_i^ρ , $1 \leq i \leq n$, $\rho \in \{0, 1\}$, so that each of these sets is a finite union of rational balls and so that the closures \overline{W}_i^0 and \overline{W}_i^1 are disjoint and $f(A_i^\rho) \subseteq W_i^\rho$ for all $i \in \{1, \dots, n\}$ and $\rho \in \{0, 1\}$. Let $k_0 \in \mathbb{N}$ be such that $2 \cdot 2^{-k_0} < d(W_i^0, W_i^1)$ for each $i \in \{1, \dots, n\}$ (such k_0 certainly exists since \overline{W}_i^0 and \overline{W}_i^1 are compact and disjoint for each $i \in \{1, \dots, n\}$).

Using Proposition 3.7, Lemma 3.4 and Proposition 4.1 we conclude that for each $k \in \mathbb{N}$ there exists $l \in \mathbb{N}$ with the following properties:

$$\widehat{\mathcal{H}}_l \text{ is an } n\text{-chain, } \mathcal{H}_l \text{ covers } E, \partial \mathcal{H}_l \text{ covers } S, \quad (4.8)$$

$$\text{fmesh}(l) < 2^{-(k+k_0)}, \mathcal{H}_l \text{ is } 2^{-(k+k_0)} \text{ - proper} \quad (4.9)$$

and

$$\bigcup \left(\partial_i^\rho \widehat{\mathcal{H}}_l \right) \subseteq W_i^\rho \quad (4.10)$$

for all $i \in \{1, \dots, n\}$ and $\rho \in \{0, 1\}$.

As in the proof of Theorem 4.8 we conclude that there exists a computable function $g : \mathbb{N} \rightarrow \mathbb{N}$ such that (4.8), (4.9) and (4.10) hold for each $k \in \mathbb{N}$ and $l = g(k)$. Let $k \in \mathbb{N}$. By Proposition 3.9 for each $x \in \bigcup \mathcal{H}_{g(k)}$ there exists $y \in E$ such that $d(x, y) < 7 \cdot 2^{-k}$ and therefore E is computable. \square

Let us mention that a computable n -cell need not be computably homeomorphic to the unit ball in \mathbb{R}^n . It has been shown in [7] that there exists a computable arc E in \mathbb{R}^2 with computable endpoints, but such that there exists no homomorphism $[0, 1] \rightarrow E$ which is a computable function. Similarly, a computable $(n - 1)$ -sphere need not be computably homeomorphic to the unit sphere in \mathbb{R}^n ([7]).

5. CONCLUSION

In this paper we have seen that topology plays an important role regarding the computability of co-c.e. sets in computable metric space. We have seen that the topological types of an arbitrary dimensional sphere and an arbitrary dimensional cell behave well from this viewpoint not just in Euclidean space but in any computable metric space which is locally computable, in particular in any computable metric space which has the effective covering property and which is locally compact. Such a computable metric space is for example the Hilbert cube I^∞ , equipped with a natural computability structure (see e.g. [5]).

It should be mentioned that co-c.e. spheres, as well as co-c.e. cells with co-c.e. boundary spheres, need not be computable in a computable metric space which is not locally computable. Moreover, by [6], there are examples of computable metric spaces X and Y such that X has the effective covering property and Y is compact, but such that both X and Y have noncomputable co-c.e. topological circles and a noncomputable co-c.e. arcs with computable endpoints.

ACKNOWLEDGEMENTS

The author would like to thank the anonymous referees for their careful work and many useful suggestions and corrections. Furthermore, the author is grateful to Professor Sibe Mardešić and Professor Ivan Ivanišić for their helpful comments.

REFERENCES

- [1] Vasco Brattka. Plottable real number functions and the computable graph theorem. *SIAM J. Comput.*, 38(1):303–328, 2008.
- [2] Vasco Brattka and Gero Presser. Computability on subsets of metric spaces. *Theoretical Computer Science*, 305:43–76, 2003.
- [3] Charles O. Christenson and William L. Voxman. Aspects of Topology. *Marcel Dekker, Inc., New York*, 1977.
- [4] Ryszard Engelking. Dimension Theory. *PWN – Polish Scientific Publishers, Warszawa*, 1978.
- [5] Zvonko Iljazović. Chainable and Circularly Chainable Co-c.e. Sets in Computable Metric Spaces. *Journal of Universal Computer Science*, 15(6):1206–1235, 2009.
- [6] Zvonko Iljazović. Local Computability of Computable Metric Spaces and Computability of Co-c.e. Continua. To appear in *Glasnik matematički*.
- [7] Joseph S. Miller. Effectiveness for Embedded Spheres and Balls. *Electronic Notes in Theoretical Computer Science*, 66:127–138, 2002.
- [8] James R. Munkres. Elements of Algebraic Topology. *Addison-Wesley Publishing Company, Menlo Park, California*, 1984.
- [9] Ernst Specker. Der Satz vom Maximum in der rekursiven Analysis. Constructivity in Mathematics (A. Heyting, ed.). *North Holland Publ. Comp., Amsterdam*, 254–265, 1959.
- [10] Klaus Weihrauch. *Computable Analysis Springer, Berlin*, 2000.