RANDOMNESS EXTRACTION AND ASYMPTOTIC HAMMING DISTANCE

CAMERON E. FREER a AND BJØRN KJOS-HANSSEN b

- ^a Computer Science and Artificial Intelligence Laboratory, Massachusetts Institute of Technology e-mail address: freer@math.mit.edu
- b Department of Mathematics, University of Hawai'i at Mānoa e-mail address: bjoernkh@hawaii.edu

ABSTRACT. We obtain a non-implication result in the Medvedev degrees by studying sequences that are close to Martin-Löf random in asymptotic Hamming distance. Our result is that the class of stochastically bi-immune sets is not Medvedev reducible to the class of sets having complex packing dimension 1.

1. Introduction

We are interested in the extent to which an infinite binary sequence X, or equivalently a set $X \subseteq \omega$, that is algorithmically random (Martin-Löf random) remains useful as a randomness source after modifying some of the bits. Usefulness here means that some algorithm (extractor) can produce a Martin-Löf random sequence from the result Y of modifying X. For further motivation see Subsection 1.2 and Section 3.

A set that lies within a small Hamming distance of a random set may be viewed as produced by an adaptive adversary corrupting or fixing some bits after looking at the original random set. Similar problems in the finite setting have been studied going back to Ben-Or and Linial [1].

If A is a finite set and $\sigma, \tau \in \{0,1\}^A$, then the Hamming distance $d(\sigma, \tau)$ is given by

$$d(\sigma,\tau) = \left| \left\{ n : \sigma(n) \neq \tau(n) \right\} \right|.$$

2012 ACM CCS: [Theory of computation]: Models of computation—Computability / Probabilistic computation; Computational complexity and cryptography—Problems, reductions and completeness; Randomness, geometry and discrete structures—Pseudorandomness and derandomization.

2010 Mathematics Subject Classification: 03D30, 03D32, 68Q30.

Key words and phrases: randomness extraction, algorithmic randomness, Hamming distance.

Submitted Dec. 11, 2012

Published

Sep. 25, 2013

^a Work on this publication was made possible through the support of a grant from the John Templeton Foundation. The opinions expressed in this publication are those of the authors and do not necessarily reflect the views of the John Templeton Foundation.

^b This material is based upon work supported by the National Science Foundation under Grant No. 0901020.

Let the collection of all infinite computable subsets of ω be denoted by \mathfrak{C} . Let $p:\omega\to\omega$. For $X,Y\in 2^\omega$ and $N\subseteq\omega$ we define a notion of proximity, or similarity, by

$$X \sim_{p,N} Y \iff (\exists n_0)(\forall n \in N, n \ge n_0)(d(X \upharpoonright n, Y \upharpoonright n) \le p(n)).$$

We will study the effective dimension of sequences that are $\sim_{p,N}$ to certain algorithmically random reals for suitably slow-growing functions p.

We use the following notation for a kind of neighborhood around X.

$$[X]_{p,N} = \{Y : Y \sim_{p,N} X\}.$$

Moreover, for a collection \mathcal{A} of subsets of ω ,

$$[\mathcal{A}]_{p,N} = \bigcup \{ [X]_{p,N} : X \in \mathcal{A} \}.$$

Turing functionals as random variables. Since a random variable must be defined for all elements of the sample space, we consider a Turing functional Φ to be a map into

$$\Omega := 2^{<\omega} \cup 2^{\omega}.$$

Setting the domain of Φ to also be Ω allows for composing maps. Let

$$\Lambda^X(n) = X(n).$$

Thus $\Lambda: 2^{\omega} \to 2^{\omega}$ is the identity Turing functional.

We define a probability measure λ on Ω called *Lebesgue (fair-coin) measure*, whose σ -algebra of λ -measurable sets is

$$\mathfrak{F} = \{ \mathcal{S} \subseteq \Omega : \mathcal{S} \cap 2^{\omega} \text{ is Lebesgue measurable} \},$$

by letting $\lambda(\mathcal{S})$ equal the fair-coin measure of $\mathcal{S} \cap 2^{\omega}$. Thus $\lambda(2^{\omega}) = 1$ and $\lambda(2^{<\omega}) = 0$, that is, λ is concentrated on the functions that are actually total.

The distribution of Φ is the measure $\mathcal{S} \mapsto \lambda\{X : \Phi^X \in \mathcal{S}\}$, defined on \mathfrak{F} . Thus the distribution of Λ is λ .

If $X \in 2^{\omega}$ then X is called a real, a set, or a sequence depending on context. If $I \subseteq \omega$ then $X \upharpoonright I$ denotes X, viewed as a function, restricted to the set I. We denote the cardinality of a finite set A by |A|. Regarding X,Y as subsets of ω and letting + denote sum mod two, note that $(X+Y) \cap n = \{k < n : X(k) \neq Y(k)\}$ and generally for a set $I \subseteq \omega$, $(X+Y) \cap I = \{k \in I : X(k) \neq Y(k)\}$.

For an introduction to algorithmic randomness the reader may consult the recent books by Nies [10] and Downey and Hirschfeldt [4]. Let MLR denote the set of Martin-Löf random elements of 2^{ω} . For a binary relation R we use a set-theoretic notation for image,

$$R[\![A]\!] = \{y : (\exists x \in A)(\langle x, y \rangle \in R)\}.$$

Let the use $\varphi^X(n)$ be the largest number used in the computation of $\Phi^X(n)$. We write

$$\Phi^X(n) \downarrow @s$$

if $\Phi^X(n)$ halts by stage s, with use at most s; if this statement is false, we write $\Phi^X(n) \uparrow @s$. We may assume that the running time of a Turing reduction is the same as the use, because any X-computable upper bound on the use is a reasonable notion of use.

For a set $\mathcal{A} \subseteq 2^{\omega}$, let

Interior_{p,N}(
$$\mathcal{A}$$
) = { $X : (\forall Y \sim_{p,N} X)(Y \in \mathcal{A})$ }
 $\subseteq \text{Interior}_*(\mathcal{A}) = {X : (\forall Y = X)(Y \in \mathcal{A})} \subseteq \mathcal{A}$

where =* denotes almost equality for all but finitely many inputs. It is easy to see that

$$Interior_{p,N}(MLR) = \emptyset$$

whenever $N \subseteq \omega$ and p is unbounded.

Definition 1.1 (Effective convergence). Let $\{a_n\}_{n\in\omega}$ be a sequence of real numbers.

- $\{a_n\}_{n\in\omega}$ converges to ∞ effectively if there is a computable function N such that for all k and all $n \geq N(k)$, $a_n \geq k$.
- $\{a_n\}_{n\in\omega}$ converges to 0 effectively if the sequence $\{a_n^{-1}\}_{n\in\omega}$ converges to ∞ effectively.

Definition 1.2. For a sequence of real numbers $\{a_n\}_{n\in\omega}$, $\lim_{n\to\infty}^* a_n$ is the real number to which a_n converges effectively, if any; and is undefined if no such number exists.

As a kind of effective big-O notation, $p_n = \omega^*(q_n)$ means $\lim_{n\to\infty}^* q_n/p_n = 0$, i.e., q_n/p_n goes to zero effectively.

Central Limit Theorem. Let \mathcal{N} be the cumulative distribution function for a standard normal random variable; so

$$\mathcal{N}(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{x} e^{-t^2/2} dt.$$

Let \mathbb{P} denote fair-coin probability on Ω . We may write

$$\mathbb{P}(\text{Event}) = \mathbb{P}(\{X : X \in \text{Event}\}) = \lambda \{X : X \in \text{Event}\}.$$

We will make use of the following quantitative version of the central limit theorem.

Theorem 1.3 (Berry-Esséen¹). Let $\{X_n\}_{n\geq 1}$ be independent and identically distributed real-valued random variables with the expectations $\mathbb{E}(X_n) = 0$, $\mathbb{E}(X_n^2) = \sigma^2$, and $\mathbb{E}(|X_n|^3) = \rho < \infty$. Then there is a constant d (with $.41 \leq d \leq .71$) such that for all x and n,

$$\left| \mathbb{P}\left(\frac{\sum_{i=1}^{n} X_i}{\sigma \sqrt{n}} \le x \right) - \mathcal{N}(x) \right| \le \frac{d\rho}{\sigma^3 \sqrt{n}}.$$

We are mostly interested in the case $X_n = X(n) - \frac{1}{2}$, $X(n) \in \{0, 1\}$, for $X \in 2^{\omega}$ under λ , in which case $\sigma = 1/2$.

1.1. New Medvedev degrees. Let \leq_s denote Medvedev (strong) reducibility and let \leq_w denote Muchnik (weak) reducibility. A recent survey of the theory behind these reducibilities is Hinman [8].

Definition 1.4 (see, e.g., [9]). A set X is *immune* if for each $N \in \mathfrak{C}$, $N \not\subseteq X$. If $\omega \setminus X$ is immune then X is *co-immune*. If X is both immune and co-immune then X is *bi-immune*.

Definition 1.5. A set X is *stochastically bi-immune* if for each set $N \in \mathfrak{C}$, $X \upharpoonright N$ satisfies the strong law of large numbers, i.e.,

$$\lim_{n\to\infty}\frac{|X\cap N\cap n|}{|N\cap n|}=\frac{1}{2}.$$

¹See for example Durrett [5].

Definition 1.6. Let $0 \le \mathfrak{p} < 1$. A sequence $X \in 2^{\omega}$ is \mathfrak{p} -stochastically dominated if for each $L \in \mathfrak{C}$,

$$\limsup_{n\to\infty}\frac{|L\cap n|}{n}>0\quad\Longrightarrow\quad (\exists M\in\mathfrak{C})\quad M\subseteq L\quad\text{and}\quad \limsup_{n\to\infty}\frac{|X\cap M\cap n|}{|M\cap n|}\leq\mathfrak{p}.$$

The class of stochastically dominated sequences is denoted $SD = SD_{\mathfrak{p}}$. If $\omega \setminus X \in SD_{\mathfrak{p}}$ then we write $X \in SD^{\mathfrak{p}}$ and say that X is stochastically dominating.

Let IM denote the set of immune sets, CIM the set of co-immune sets, and W3R the set of weakly 3-random sets. Let K denote prefix-free Kolmogorov complexity.

Definition 1.7 (see, e.g., [4, Ch. 13]). The effective Hausdorff dimension of $A \in 2^{\omega}$ is

$$\dim_H(A) = \liminf_{n \in \omega} \frac{K(A \upharpoonright n)}{n}.$$

The complex packing dimension of $A \in 2^{\omega}$ is

$$\dim_{cp}(A) = \sup_{N \in \mathfrak{C}} \inf_{n \in N} \frac{K(A \upharpoonright n)}{n}.$$

The effective packing dimension of $A \in 2^{\omega}$ is

$$\dim_p(A) = \limsup_{n \in \omega} \frac{K(A \upharpoonright n)}{n}.$$

Proposition 1.8. For all $A \in 2^{\omega}$,

$$0 \le \dim_H(A) \le \dim_{cp}(A) \le \dim_p(A) \le 1.$$

Proof. The inequality $\dim_H(A) \leq \dim_{cp}(A)$ uses the fact that each cofinite set $N \subseteq \omega$ is in \mathfrak{C} . The inequality $\dim_{cp}(A) \leq \dim_p(A)$ uses the fact that each $N \in \mathfrak{C}$ is an infinite subset of ω .

By examining the complex packing dimension of reals that are $\sim_{p,N}$ to a Martin-Löf random real for p growing more slowly than $n/(\log n)$, we will derive our main result, which states the existence, for each Turing reduction Φ , of a set Y of complex packing dimension 1 for which Φ^Y is not stochastically bi-immune.

1.2. Relation of our results to other recent results. Jockusch and Lewis [9] prove that the class of bi-immune sets is Medvedev reducible to the class of almost diagonally non-computable functions DNC*, i.e., functions f such that $f(x) = \varphi_x(x)$ for at most finitely many x. Downey, Greenberg, Jockusch, and Milans [3] show that DNC₃ (the class of DNC functions taking values in $\{0,1,2\}$) and hence also its superset DNC*, is not Medvedev above the class of Kurtz random sets. We do not know whether the class of stochastically bi-immune sets is Medvedev reducible to the class of DNC* functions. We show in Theorem 4.3 below that from a set of complex packing dimension 1 one cannot uniformly compute a stochastically bi-immune set; on the other hand, to compute a DNC* function from a set of complex packing dimension 1 one would apparently also need to know the witnessing set $N \in \mathfrak{C}$.

Definition 1.9 (see, e.g., [10, Def. 7.6.4]). A sequence $X \in 2^{\omega}$ is Mises-Wald-Church (MWC) stochastic if no partial computable monotonic selection rule can select a biased subsequence of X, i.e., a subsequence where the relative frequencies of 0s and 1s do not converge to 1/2.




Figure 1: Some Medvedev degrees. The fact that (cp, 1) is not Medvedev above SBI is Theorem 4.3.

→ Included in

~~> Not Medvedev above

→ Medvedev above

Figure 2: Meaning of arrows.

Definition 1.10. A sequence $X \in 2^{\omega}$ is BI² (bi-immune for sets of size two) if there is no computable collection of disjoint finite sets of size 2 on which the set omits a certain pattern such as 01. More precisely, X is BI² if for each computable disjoint collection $\{T_n : n \in \omega\}$ where each T_n has cardinality two, say $T_n = \{s_n, t_n\}$ where $s_n < t_n$, and each $P \subseteq \{0, 1\}$, there is an n such $X(s_n) = P(0)$ and $X(t_n) = P(1)$.

Each von Mises-Wald-Church stochastic (MWC-stochastic) set is stochastically biimmune. Our main theorem implies that a set of complex packing dimension 1 does not necessarily uniformly compute a MWC-stochastic set. This consequence is not really new with the present paper, however, because the fact that DNC₃ is not Medvedev above BI² is implicit in Downey, Greenberg, Jockusch, and Milans [3] as pointed out to us by Joe Miller. The situation is diagrammatically illustrated in Figure 1, with notation defined in Figures 2 and 3. In the future we could hope to replace complex packing dimension by effective Hausdorff dimension in Theorem 4.3.

Abbreviation	Unabbreviation	Definition
DNC_n	Diagonally non-computable function in n^{ω}	
MLR	Martin-Löf random	
KR	Kurtz random (weakly 1-random)	
MWC	Mises-Wald-Church stochastic	1.9
SBI	Stochastically bi-immune	1.5
BI	bi-immune	1.4
BI^2	bi-immune for sets of size two	1.10
(H,1)	effective Hausdorff dimension 1	1.7
(cp,1)	complex packing dimension 1	1.7

Figure 3: Abbreviations used in Figure 1.

2. Hamming space

The Hamming distance between a point and a set of points is defined by $d(y, A) := \min_{a \in A} d(y, a)$. The r-neighborhood of a set $A \subseteq \{0, 1\}^n$ is

$$\Gamma_r(A) = \{ y \in \{0,1\}^n : d(y,A) \le r \}.$$

In particular,

$$\Gamma_r(\{c\}) = \{y \in \{0,1\}^n : d(y,c) \le r\},\$$

and

$$\Gamma_r(A) = \bigcup_{a \in A} \Gamma_r(\{a\}).$$

A Hamming-sphere² with center $c \in \{0,1\}^n$ is a set $S \subseteq \{0,1\}^n$ such that for some k,

$$\Gamma_k(\{c\}) \subseteq S \subseteq \Gamma_{k+1}(\{c\}).$$

Theorem 2.1 (Harper [7]; see also Frankl and Füredi [6]). For each $n, r \ge 1$ and each set $A \subseteq \{0,1\}^n$, there is a Hamming-sphere $S \subseteq \{0,1\}^n$ such that

$$|A| = |S|$$
, and $|\Gamma_r(A)| \ge |\Gamma_r(S)|$.

Following Buhrman et al. [2], we write

$$b(n,k) := \binom{n}{0} + \dots + \binom{n}{k}.$$

Note that for all $c \in \{0,1\}^n$, $|\Gamma_k(\{c\})| = b(n,k)$

If the domain of σ is an interval I in ω rather than an initial segment of ω , we may emphasize I by writing

$$B_r^I(\sigma) = \Gamma_r(\{\sigma\}) = \{\tau \in \{0,1\}^I : d(\sigma,\tau) \le r\}.$$

 $\mathbb P$ denotes the uniform distribution on $\{0,1\}^I,$ so by definition

$$\mathbb{P}(E) = \frac{|E|}{2^{|I|}}.$$

Recall that D_m is the m^{th} canonical finite set. The intuitive content of Lemma 2.2 below is that a medium size set is unlikely to contain a random large ball. (Note that we do not assume the sets I_m are disjoint.)

²A Hamming-sphere is more like a ball than a sphere, but the terminology is entrenched.

Lemma 2.2. Let $\chi \in \omega^{\omega}$. Suppose

$$\lim_{n \to \infty}^{*} \chi(n) / \sqrt{n} = \infty. \tag{2.1}$$

Let $f \in \omega^{\omega}$ be a computable function. Let $I_m = D_{f(m)}$ and $n_m = |I_m|$. Suppose

$$\lim_{m \to \infty}^* n_m = \infty. \tag{2.2}$$

For each $m \in \omega$ let $E_m \subseteq \{0,1\}^{I_m}$. Suppose $\limsup_{m \to \infty} \mathbb{P}(E_m) \leq \mathfrak{p}$ where $0 < \mathfrak{p} < 1$ is computable. Writing

$$B_{\chi(n)}(X)$$
 for $B_{\chi(n_m)}^{I_m}(X \upharpoonright I_m)$,

we have

$$\lim_{m \to \infty}^* \mathbb{P}(\{X : B_{\chi(n)}(X) \subseteq E_m\}) = 0. \tag{2.3}$$

Moreover, for each $m_0 \in \omega$ and computable $\mathfrak{q} \in (\mathfrak{p}, 1)$ there is a modulus of effective convergence in (2.3) that works for all sets $\{E_m\}_{m \in \omega}$ such that for all $m \geq m_0$, $\mathbb{P}(E_m) \leq \mathfrak{q}$.

Proof. Let $X \in 2^{\omega}$ be a random variable with $X =_d \Lambda$, and

$$S^{(m)} = \sum_{i \in I_m} X(i).$$

Let

$$f_m(x) = \mathbb{P}\left(\frac{S^{(m)} - n/2}{\sqrt{n}/2} \le x\right)$$

We have 3

$$|f_m(x) - \mathcal{N}(x)| \le \frac{d}{\sqrt{n_m}}.$$

Since $\lim_{m\to\infty}^* n_m = \infty$, $\lim_{m\to\infty}^* \frac{d}{\sqrt{n_m}} = 0$. So

$$\lim_{m \to \infty} \sup_{x} |f_m(x) - \mathcal{N}(x)| = 0.$$
 (2.4)

Let $r = r_m$ be such that

$$b(n,r) \le |E_m| < b(n,r+1).$$

Let

$$a_m = \frac{r_m - \frac{n}{2}}{\sqrt{n/2}},$$

and let

$$b_m = a_m + \frac{1}{\sqrt{n}/2} - \frac{\chi(n)}{\sqrt{n}/2}.$$

By (2.4),

$$\lim_{m \to \infty}^* |f_m(b_m) - \mathcal{N}(b_m)| = 0.$$
(2.5)

$$\left| \mathbb{P} \left(\frac{\sum_{i \in I_m} Y_i}{\sigma \sqrt{n}} \le x \right) - \mathcal{N}(x) \right| \le \frac{d\rho}{\sigma^3 \sqrt{n}} = \frac{d}{\sqrt{n}},$$

where $\rho = 1/8 = \mathbb{E}(|Y_i|^3)$, and $\sigma = 1/2$ is the standard deviation of X_i (and Y_i).

³Indeed, let $Y_i = X_i - \mathbb{E}(X_i)$ where $\mathbb{E}(X_i) = \frac{1}{2}$ is the expected value of X_i , so $\mathbb{E}(Y_i) = 0$. By the Berry-Esséen Theorem 1.3, for all x

We have

$$\lim_{m \to \infty} \sup f_m(a_m) = \lim_{m \to \infty} \mathbb{P}\left(\frac{S^{(m)} - n/2}{\sqrt{n}/2} \le \frac{r_m - \frac{n}{2}}{\sqrt{n}/2}\right)$$
$$= \lim_{m \to \infty} \sup \mathbb{P}(S^{(m)} \le r_m) = \lim_{m \to \infty} \sup \frac{b(n, r)}{2^n} \le \lim_{m \to \infty} \sup \mathbb{P}(E_m) \le \mathfrak{p}.$$

Since $f_m \to \mathcal{N}$ uniformly, it follows that

$$\lim_{m\to\infty}\sup \mathcal{N}(a_m)\leq \mathfrak{p},$$

and so as \mathcal{N} is strictly increasing,

$$\limsup_{m \to \infty} a_m \le \mathcal{N}^{-1}(\mathfrak{p}) \quad (= 0 \text{ if } \mathfrak{p} = 1/2).$$

Let m_0 be such that for all $m \geq m_0$,

$$a_m + \frac{1}{\sqrt{n_m/2}} \le \mathcal{N}^{-1}(\mathfrak{p}) + 1.$$

Since by assumption $\lim_{n\to\infty}^* \chi(n)/\sqrt{n} = \infty$, we have that b_m is the sum of a term that goes effectively to $-\infty$, and a term that after m_0 never goes above $\mathcal{N}^{-1}(\mathfrak{p}) + 1$ again. Thus

$$\lim_{m\to\infty}^* b_m = -\infty.$$

It is this rate of convergence that is transformed in the rest of the proof. Now

$$\lim_{m\to\infty}^* \mathcal{N}(b_m) = 0.$$

Hence by (2.5), $\lim_{m\to\infty}^* f_m(b_m) = 0$.

Let us write

$$B_t(X) := B_t^{I_m}(X \upharpoonright I_m),$$

considering $X \upharpoonright I_m$ as a string of length n. By Harper's Theorem 2.1, we have a Hamming sphere H with

$$|H| = |\neg E_m|$$
 and $|\Gamma_{\chi(n)}(\neg E_m)| \ge |\Gamma_{\chi(n)}(H)|$.

Then

$$\mathbb{P}(\{X:X\in\Gamma_{\chi(n)}(\neg E_m)\})\geq \mathbb{P}(\{X:X\in\Gamma_{\chi(n)}(H)\}).$$

Therefore

$$\mathbb{P}(\{X:X\not\in\Gamma_{\chi(n)}(\neg E_m)\})\leq\mathbb{P}(\{X:X\not\in\Gamma_{\chi(n)}H)\}).$$

Let \widehat{H} be the complement of H. If the Hamming sphere H is centered at $c \in \{0,1\}^n$ then clearly \widehat{H} is a Hamming sphere centered at \overline{c} , where $\overline{c}(k) = 1 - c(k)$. Since

$$\left|\widehat{H}\right| = |E_m| < b(n, r+1),$$

we have $\widehat{H} \subset \Gamma_{r+1}(\{\overline{c}\})$. So we have:

$$\mathbb{P}(\{X : B_{\chi(n)}(X) \subseteq E_m\}) \leq \mathbb{P}(\{X : B_{\chi(n)}(X) \subseteq \widehat{H}\})$$

$$< \mathbb{P}(\{X : B_{\chi(n)}(X) \subseteq \Gamma_{r+1}(\{\overline{c}\})\}) = \frac{b(n, r+1-\chi(n))}{2^n}$$

$$= \mathbb{P}[S^{(m)} \leq r+1-\chi(n)] = \mathbb{P}\left[\frac{S^{(m)} - \frac{n}{2}}{\sqrt{n}/2} \leq \frac{r+1-\frac{n}{2}-\chi(n)}{\sqrt{n}/2}\right]$$

$$= f_m \left(a_m + \frac{1}{\sqrt{n}/2} - \frac{\chi(n)}{\sqrt{n}/2} \right) = f_m(b_m).$$

Since we showed that $\lim_{m\to\infty}^* f_m(b_m) = 0$, and since by assumption $\lim_{m\to\infty}^* n_m = \infty$,

$$\lim_{m \to \infty}^* \mathbb{P}(\{X : B_{\chi(n)}(X) \subseteq E_m\}) = 0.$$

3. Turing reductions that preserve randomness

The way we will obtain our main result Theorem 4.3 is by proving essentially that for any "randomness extractor" Turing reduction, and any random input oracle, a small number of changes to the oracle will cause the extractor to fail to produce a random output. This would be much easier if we restricted attention to Turing reductions having disjoint uses on distinct inputs, since we would be working with independent random variables. Indeed, one can give an easy proof in that case, which we do not include here. The main technical achievement of the present paper is to be able to work with overlapping use sets; key in that respect is Lemma 3.3 below. The number of changes to the random oracle that we need to make is small enough that the modified oracle has complex packing dimension 1. We were not able to set up the construction so as to guarantee effective Hausdorff dimension 1 (or even greater than 0); this may be an avenue for future work.

For a set of pairs E, we have the projections $E^x = \{y : (x,y) \in E\}$ and $E_y = \{x : (x,y) \in E\}$.

Lemma 3.1. Let μ_1 and μ_2 be probability measures on sample spaces Ω_1 and Ω_2 and let E be a measurable subset of $\Omega_1 \times \Omega_2$. Suppose that η , α , and δ are positive real numbers such that

$$\mu_1 E_y > \eta \quad (\forall y \in \Omega_2), \quad and$$
 (3.1)

$$\mu_1\{x: \mu_2 E^x \le \alpha\} \ge 1 - \delta. \tag{3.2}$$

Then $\eta < \alpha + \delta$.

Proof. By Fubini's theorem,

$$\eta < \int_{\Omega_2} \mu_1(E_y) d\mu_2(y) = \iint_{\Omega_1 \times \Omega_2} E(x, y) d\mu_1(x) d\mu_2(y) = \int_{\Omega_1} \mu_2(E^x) d\mu_1(x)
\leq \alpha \cdot \mu_1 \{x : \mu_2(E^x) \leq \alpha\} + 1 \cdot \mu_1 \{x : \mu_2(E^x) \geq \alpha\} \leq \alpha \cdot 1 + 1 \cdot \delta. \qquad \square$$

Definition 3.2. For a real X and a string σ of length n,

$$(\sigma \searrow X)(n) = \begin{cases} \sigma(n) & \text{if } n < |\sigma|, \\ X(n) & \text{otherwise,} \end{cases}$$

and

$$(\sigma^{\widehat{}}X)(n) = \begin{cases} \sigma(n) & \text{if } n < |\sigma|, \\ X(n - |\sigma|) & \text{otherwise.} \end{cases}$$

Thinking of σ and X as functions we may write

$$\sigma \searrow X = \sigma \cup (X \upharpoonright \omega \backslash |\sigma|)$$

and thinking in terms of concatenation we may write

$$\sigma^{\hat{}}X = \sigma X.$$

Lemma 3.3. Let Φ be a Turing reduction such that

$$\lambda(\Phi^{-1}[SD_{\mathfrak{p}}]) = 1 \tag{3.3}$$

and let $\Phi_{\sigma}^{X} = \Phi^{\sigma \searrow X}$. Then for any finite set $\Sigma \subseteq 2^{<\omega}$,

$$(\forall \varepsilon > 0)(\forall i_0)(\exists i > i_0)(\forall \sigma \in \Sigma)$$

$$\mathbb{P}(\{X \mid \Phi_{\sigma}^{X}(i) = 1\}) \le \mathfrak{p} + \varepsilon.$$

Proof. First note that for all $\sigma \in 2^{<\omega}$, $\lambda(\Phi_{\sigma}^{-1}[SD_{\mathfrak{p}}] = 1$ as well.

Suppose otherwise, and fix ε , i_0 and Σ such that

$$(\forall i > i_0)(\exists \sigma \in \Sigma) \quad \mathbb{P}(\Phi_{\sigma}(i) = 1) > \mathfrak{p} + \varepsilon.$$

By density of the rationals in the reals we may assume ε is rational and hence computable. Since there are infinitely many i but only finitely many σ , it follows that there is some σ such that

$$(\exists^{\infty} k > i_0) \quad \mathbb{P}(\Phi_{\sigma}(k) = 1) > \mathfrak{p} + \varepsilon \tag{3.4}$$

and in fact

$$\limsup |\{k < n : \mathbb{P}(\Phi_{\sigma}(k) = 1) > \mathfrak{p} + \varepsilon\}| / n > 0.$$

Fix such a σ and let $\Psi = \Phi_{\sigma}$. Let $\{\ell_n\}_{n \in \omega}$ be infinitely many values of k in (3.4) listed in increasing order; note that $L = \{\ell_n\}_{n \in \omega}$ may be chosen as a computable sequence.

For an as yet unspecified subsequence $\mathcal{K} = \{k_n\}_{n \in \omega}, \ \mathcal{K} \subseteq L$, let

$$E = \{(X, n) : \Psi^X(k_n) = 1\}. \tag{3.5}$$

We obtain then also projections $E_n = \{X : \Psi^X(k_n) = 1\}, E^X = \{n : \Psi^X(k_n) = 1\}.$ By (3.4) we have for all $n \in \omega$,

$$\lambda E_n > \mathfrak{p} + \varepsilon. \tag{3.6}$$

The fraction of events E_n that occur in $N = \{0, \dots, N-1\}$ for X is denoted

$$e_N^X = \frac{\left| E^X \cap N \right|}{N}$$

By assumption (3.3),

$$\lambda \left\{ X : (\exists \mathcal{K} \subseteq L)(\exists M)(\forall N \ge M) \left(e_N^X \le \mathfrak{p} + \frac{\varepsilon}{2} \right) \right\} = 1.$$

Thus there is an M and a K (using that \mathfrak{C} is countable) such that

$$\lambda \left\{ X : e_M^X \le \mathfrak{p} + \frac{\varepsilon}{2} \right\} \ge \lambda \left\{ X : (\forall N \ge M) e_N^X \le \mathfrak{p} + \frac{\varepsilon}{2} \right\} \ge 1 - \frac{\varepsilon}{3}. \tag{3.7}$$

Let Ω_1 be the unit interval [0,1]. Let $\Omega_2 = M = \{0,1,\ldots,M-1\}$. Let $\mu_1 = \lambda$. Let $\mu_2 = \text{card}$ be the counting measure on the finite set $M = \{0,1,\ldots,M-1\}$, so that for a finite set $A \subset M$, $\operatorname{card}(A)$ is the cardinality of A. Let $\eta = \mathfrak{p} + \varepsilon$, $\alpha = \mathfrak{p} + \varepsilon/2$, and $\delta = \varepsilon/3$, and note that $\eta > \alpha + \delta$. By (3.6), (3.7) and Lemma 3.1, $\eta < \alpha + \delta$, a contradiction.

⁴In this case, $\int \mu_1(E_y) d\mu_2(y) = \int \mu_1(E_n) d\mu_2(n) = \int \lambda(E_n) d \operatorname{card}(n) = \sum_{n \in \Omega_2} \lambda(E_n) \operatorname{card}(\{n\}) = \sum_{n=0}^{M-1} \lambda(E_n) \cdot 1.$

4. Extraction and Hamming distance

Theorem 4.1. Let $\mathfrak{p} < 1$ be computable. Let $p : \omega \to \omega$ be any computable function such that $p(n) = \omega^*(\sqrt{n})$. Let Φ be a Turing reduction. There exists an $N \in \mathfrak{C}$ and an almost sure event A such that

$$\mathcal{A} \cap \operatorname{Interior}_{p,N}(\Phi^{-1}[\![\mathcal{A}]\!]) = \emptyset$$

Proof. Let

$$\mathcal{A} := \mathrm{W3R} \subset \mathrm{MLR} \subset \mathrm{CIM} \cap \mathrm{IM} \cap (\mathrm{SD}_{\mathfrak{p}} \cup \mathrm{SD}^{\mathfrak{p}}).$$

We show

(1) If
$$\lambda(\Phi^{-1}[SD]_{\mathfrak{p}}) = 1$$
, or $\lambda(\Phi^{-1}[SD^{\mathfrak{p}}]) = 1$, then
$$MLR \cap Interior_{\mathfrak{p},N}(\Phi^{-1}[CIM]) = \varnothing,$$

$$MLR \cap Interior_{p,N}(\Phi^{-1}\llbracket IM \rrbracket) = \emptyset$$
, respectively.

(2) Otherwise; then

$$W3R \cap Interior_*(\Phi^{-1}[SD_{\mathfrak{p}}]) = \varnothing$$

and

$$W3R \cap Interior_*(\Phi^{-1} \llbracket SD^{\mathfrak{p}} \rrbracket) = \varnothing$$

Proof of (2): If we are not in case (1) then $\lambda\{X \mid \Phi^X \in \mathrm{SD}_{\mathfrak{p}}\} < 1$, so by the 0-1 Law, $\lambda\{X \mid (\forall Y = X)(\Phi^Y \in \mathrm{SD}_{\mathfrak{p}})\} = 0$. This is (contained in) a Π_3^0 null class, so if $X \in \mathrm{W3R}$ then $(\exists Y = X)(\Phi^Y \notin \mathrm{SD}_{\mathfrak{p}})$ hence we are done.

Proof of (1): By Lemma 3.3,

$$(\exists \mathfrak{p} < 1)(\forall \varepsilon > 0)(\forall n)(\forall i)(\exists i' > i)(\forall \sigma \in 2^{=n})$$

$$\mathbb{P}(\{Z : \Phi^{\sigma \searrow Z}(i') = 1\}) \le \mathfrak{p} + \varepsilon; \tag{4.1}$$

Since Φ is total for almost all oracles, it is clear that i' is a computable function f(k,n) of $\varepsilon = 1/k$ and n. Let $g: \omega \to \omega$ be the computable function with $\lim_{n \to \infty} g(n) = \infty$ given by g(s) = 2s. Let $n_0 = 0$ and $i_0 = 0$. Assuming $s \ge 0$ and n_s and i_s have been defined, let

$$i_{s+1} = f(g(s), n_s),$$

and let n_{s+1} be large enough that

$$(\forall \sigma \in 2^{=n_s}) \quad \lambda\{Z \mid \Phi^{\sigma \searrow Z}(i_{s+1}) \uparrow @n_{s+1}\} \le \frac{1}{2s},\tag{4.2}$$

$$\lim_{s \to \infty}^* \frac{p(n_{s+1} - n_s)}{\sqrt{n_{s+1} - n_s}} = \infty, \quad \text{and} \quad \sum_{k=0}^s p(n_{k+1} - n_k) \le p(n_{s+1}).$$

Note that since i' > i in (4.1), we have $i_{s+1} > i_s$ and hence $R := \{i_0, i_1, \ldots\}$ is a computable infinite set. We now have

$$(\forall s)(\forall \sigma \in 2^{=n_s}) \quad \mathbb{P}(\{Z : \Phi^{\sigma \searrow Z}(i_{s+1}) = 1\}) \le \mathfrak{p} + \frac{1}{2s}$$

$$(4.3)$$

so

$$\mathbb{P}(\{Z:\Phi^{\sigma \searrow Z}(i_{s+1}) \downarrow = 1@n_{s+1}\}) \le \mathfrak{p} + \frac{1}{2s}.$$
(4.4)

Note $[a,b) = b \setminus a$.

Let $X \in MLR$. We aim to define $Y \sim_p X$ such that $\Phi^Y \notin MLR$. We will in fact make $Y \leq_T X$, so we define a reduction Ξ and let $Y = \Xi^X$. Since we are defining Y by modifying bits of X, the use of Ξ will be the identity function: $\xi^X(n) = n$.

Since $n_0 = 0$, $Y \upharpoonright n_0$ is the empty string. Suppose $s \ge 0$ and $Y_{\upharpoonright n_s}$ has already been defined. The set of "good" strings now is

$$\mathcal{G} = \{ \tau \succ Y_{\upharpoonright n_s} \mid \Phi^{\tau \upharpoonright n_{s+1}}(i_{s+1}) = 0 \}.$$

Define the "cost" of τ to be the additional Hamming distance to X, i.e.,

$$d(\tau) = |(X + \tau) \cap [n_s, n_{s+1})|.$$

Case 1: $\mathcal{G} \neq \emptyset$. Then let $Y_{\upharpoonright n_{s+1}}$ be any $\tau_0 \in \mathcal{G}$ of length n_{s+1} and of minimal cost, i.e., such that $d(\tau_0) = \min\{d(\tau) \mid \tau \in \mathcal{G}\}$. That is, let

$$Y_{\upharpoonright n_{s+1}} \in \operatorname*{arg\,min}_{\tau \in \mathcal{G}} d(\tau).$$

Case 2: Otherwise. Then make no further changes to X up to length n_{s+1} , i.e., let $Y_{\lceil n_{s+1} \rceil} = Y_{\lceil n_s \rceil} \setminus X_{\lceil n_{s+1} \rceil}$.

This completes the definition of Ξ and hence of Y. It remains to show that $\Phi^Y \notin MLR$. For any string σ of length n_s let

$$E_{s+1}^{\sigma} = \left\{ Z \in \{0,1\}^{[n_s,n_{s+1})} : \neg \left(\Phi^{\sigma \searrow Z}(i_{s+1}) \downarrow = 0@n_{s+1} \right) \right\}$$
$$= \left\{ Z : \Phi^{\sigma \searrow Z}(i_{s+1}) \downarrow = 1@n_{s+1} \right\}$$
$$\cup \left\{ Z : \Phi^{\sigma \searrow Z}(i_{s+1}) \uparrow @n_{s+1} \right\}$$

Since (4.2) and (4.4) hold for all strings of length n_s , in particular they hold for $\sigma = \Xi^X \upharpoonright n_s$, so

$$(\forall s) \quad \mathbb{P}(E_{s+1}^{\sigma}) \le \mathfrak{p} + \frac{1}{2s} + \frac{1}{2s} = \mathfrak{p} + \frac{1}{s}, \quad \text{hence} \quad \limsup_{s \to \infty} \mathbb{P}(E_{s+1}^{X \upharpoonright n_s}) \le \mathfrak{p}. \tag{4.5}$$

Let

$$U_s^{X \upharpoonright n_s} = \{ Z \in \{0, 1\}^{[n_s, n_{s+1})} : B_{p(n_{s+1} - n_s)}^{[n_s, n_{s+1})}(Z) \subseteq E_{s+1}^{X \upharpoonright n_s} \}.$$

Since

$$\frac{p(n_{s+1}-n_s)}{\sqrt{n_{s+1}-n_s}} \to^* \infty.$$

we can apply Lemma 2.2 and there is h(s) with $\lim_{s\to\infty}^* h(s) = 0$ and

$$\mathbb{P}(U_s^{X \upharpoonright n_s}) \le h(s)$$

that only depends on an upper bound for an s_0 such that for all $s \geq s_0$, $\mathbb{P}(E_{s+1}) \leq \mathfrak{q}$ (where $\mathfrak{p} < \mathfrak{q} < 1$ and \mathfrak{q} is just some fixed computable number). Since by (4.5) such an upper bound can be given that works for all X, actually h(s) may be chosen to not depend on X. Let

$$V_s = \{ Z : Z \in U_s^{Z \upharpoonright n_s} \},$$

then V_s is uniformly Δ_1^0 . To find the probability of V_s we note that for each of the 2^{n_s} possible beginnings of Z, there are at most $(h(s) \cdot 2^{n_{s+1}-n_s})$ continuations of Z on $[n_s, n_{s+1})$ that make $Z \in V_s$; so we compute

$$\mathbb{P}(V_s) = |\{Z \in \{0,1\}^{n_{s+1}} : Z \upharpoonright [n_s, n_{s+1}) \in U^{Z \upharpoonright n_s}\}| 2^{-n_{s+1}}$$

$$< 2^{n_s} (h(s) \cdot 2^{n_{s+1} - n_s}) 2^{-n_{s+1}} = h(s)$$

so since $\lim_{s\to\infty}^* h(s) = 0$, $\{V_s\}_{s\in\omega}$ is a Kurtz randomness test. Let $\{m_s\}_{s\in\omega}$ be a computable sequence such that $\sum_{s\geq t} h(m_s) \leq 2^{-t}$. Let $W_t = \bigcup_{s\geq t} V_{m_s}$. Then $\mathbb{P}(W_t) \leq 2^{-t}$ and W_t is uniformly Σ_1^0 and hence it is a Martin-Löf randomness test. Since $X \in MLR$,

 $X \notin W_t$ for some t and hence $X \notin V_{m_s}$ for all but finitely many s. So $\Phi^Y(m_s) = 0$ for all but finitely many s, hence $\Phi^Y \notin \text{CIM}$.

By construction, we have

$$|(X+Y)\cap[n_s,n_{s+1})| < p(n_{s+1}-n_s)$$

for all but finitely many n. Therefore

$$|(X+Y)\cap[0,n_{s+1})| \le \sum_{k=0}^{s} p(n_{k+1}-n_k) \le p(n_{s+1})$$

so $X \sim_{p,N} Y$ where $N = \{n_s : s \in \omega\}$.

4.1. Main result.

Lemma 4.2. Let $p(n) = o(\frac{n}{\log n})$ and let $N \in \mathfrak{C}$. If $X \in MLR$ and $X \sim_{p,N} Y$ then $\dim_{cp}(Y) = 1$.

Proof. Suppose there are at most p(n) many bits changed to go from $X \upharpoonright n$ to $Y \upharpoonright n$, in positions $a_1, \ldots, a_{p(n)}$. (In case there are fewer than p(n) changed bits, we can repeat a_i representing the bit 0 which we may assume is changed.) Let $(Y \upharpoonright n)^*$ be a shortest description of $Y \upharpoonright n$. From the code

$$0^{|K(Y \upharpoonright n)|} \cap 1 \cap K(Y \upharpoonright n) \cap (Y \upharpoonright n)^* \cap a_1 \cdots a_{p(n)}$$

we can effectively recover $X \upharpoonright n$. Thus

$$n - c_1 \le K(X \upharpoonright n) \le 2\log[K(Y \upharpoonright n)] + 1 + K(Y \upharpoonright n) + p(n)\log n + c_2$$

 $\le 2\log[n + 2\log n + c_3] + 1 + K(Y \upharpoonright n) + p(n)\log n + c_2.$

Hence

$$n \le {}^+ 3 \log n + K(Y \upharpoonright n) + p(n) \log n$$
, and $n - (p(n) + 3) \log n \le {}^+ K(Y \upharpoonright n)$.

Theorem 4.3. For each Turing reduction procedure Φ there is a set Y with $\dim_{cp}(Y) = 1$ such that Φ^Y is not stochastically bi-immune.

Proof. Let $p(n) = n^{2/3}$, so that $p(n) = o(n/\log n)$ and $p(n) = \omega^*(\sqrt{n})$. By the proof of Theorem 4.1 and since the sequence of numbers n_s is computable, for each weakly 3-random set X there is a set $Y \sim_{p,N} X$ (for some $N \in \mathfrak{C}$) such that Φ^Y is not both co-immune and in $\mathrm{SD}_{1/2}$, in particular $\Phi^Y \notin \mathrm{SBI}$. By Lemma 4.2, each such Y has complex packing dimension 1.

References

- Michael Ben-Or and Nati Linial, Collective coin flipping, Randomness and Computation (S. Micali, ed.), Academic Press, New York, 1989, pp. 91-115. ↑1
- [2] Harry Buhrman, Lance Fortnow, Ilan Newman, and Nikolai Vereshchagin, Increasing Kolmogorov complexity, STACS 2005, Lecture Notes in Comput. Sci., vol. 3404, Springer, Berlin, 2005, pp. 412–421. MR2151636 (2006c:68091) ↑6
- [3] Rodney G. Downey, Noam Greenberg, Carl G. Jockusch Jr., and Kevin G. Milans, *Binary subtrees with few labeled paths*, Combinatorica **31** (2011), no. 3, 285–303, DOI 10.1007/s00493-011-2634-3. MR2835294 (2012i:05336) ↑4, 5
- [4] Rodney G. Downey and Denis R. Hirschfeldt, *Algorithmic randomness and complexity*, Theory and Applications of Computability, Springer, New York, 2010. MR2732288 (2012g:03001) †2, 4
- [5] Rick Durrett, Probability: theory and examples, 4th ed., Cambridge Series in Statistical and Probabilistic Mathematics, Cambridge University Press, Cambridge, 2010. MR2722836 (2011e:60001) ↑3
- [6] Peter Frankl and Zoltán Füredi, A short proof for a theorem of Harper about Hamming-spheres, Discrete Math. **34** (1981), no. 3, 311–313, DOI 10.1016/0012-365X(81)90009-1. MR613409 (83a:05004) ↑6
- [7] Lawrence H. Harper, Optimal numberings and isoperimetric problems on graphs, J. Combinatorial Theory 1 (1966), 385–393. MR0200192 (34 #91) \uparrow 6
- [8] Peter G. Hinman, A survey of Mučnik and Medvedev degrees, Bull. Symbolic Logic 18 (2012), no. 2, 161–229, DOI 10.2178/bsl/1333560805. MR2931672 \uparrow 3
- [9] Carl G. Jockusch Jr. and Andrew E.M. Lewis, Diagonally non-computable functions and bi-immunity,
 J. Symbolic Logic. to appear. ↑3, 4
- [10] André Nies, Computability and randomness, Oxford Logic Guides, vol. 51, Oxford University Press, Oxford, 2009. MR2548883 ↑2, 4